

Klickitat Complex - Fall Chinook

This page is intentionally blank.

HATCHERY AND GENETIC MANAGEMENT PLAN
(HGMP)
Final DRAFT

Hatchery Program	Klickitat Complex - Fall Chinook
Species or Hatchery Stock	Chinook Salmon (<i>Oncorhynchus tshawytscha</i>)
Agency/Operator	Yakama Nation
Watershed and Region	Klickitat Subbasin/Columbia Gorge Province
Date Submitted	November 12, 2013
Date Last Updated	August 2013

Section 1: General Program Description

1.1 Name of hatchery or program.

Klickitat Fall Chinook – Mid-Columbia Bright

1.2 Species and population (or stock) under propagation, and ESA status.

Mid-Columbia Bright Fall Chinook Salmon (*Oncorhynchus tshawytscha*)

ESA Status: Not listed and not a candidate for listing

1.3 Responsible organization and individuals.

<i>Name (and title):</i>	Jason Rau (Complex Manager)
	Bill Sharp (YKFP Klickitat Coordinator)
<i>Agency or Tribe:</i>	Yakama Nation
<i>Address:</i>	PO Box 151 Toppenish WA 98948
<i>Telephone:</i>	(509) 865-5121
<i>Fax:</i>	(509) 865-6293
<i>Email:</i>	jayrau@ykfp.org sharp@yakama.com

Other agencies, Tribes, co-operators, or organizations involved, including contractors, and extent of involvement in the program.

Co-operators	Role
<i>Washington Dept. Fish & Wildlife (WDFW)</i>	Fish Hatchery Specialist 1
<i>United States Fish and Wildlife Service (USFWS)</i>	Health Technician and Pathologist

1.4 Funding source, staffing level, and annual hatchery program operational costs.

Funding Sources
Currently Mitchell Act. Other funding sources may include John Day Mitigation or BPA

Operational Information	Number
Full time equivalent staff	2 FTE, 3 seasonal
Estimated Annual operating cost (dollars)	\$300,000

1.5 Location(s) of hatchery and associated facilities.

<i>Broodstock source</i>	Klickitat River Fall Chinook
<i>Broodstock collection location (stream, Rkm, subbasin)</i>	Lyle Falls (Klickitat River Rkm 3.5), Klickitat River Hatchery (Rkm 68), USFWS Little White Salmon National Fish Hatchery
<i>Adult holding location (stream, Rkm, subbasin)</i>	Klickitat River Hatchery (Rkm 68), USFWS Little White Salmon National Fish Hatchery
<i>Spawning location (stream, Rkm, subbasin)</i>	Klickitat River Hatchery (Rkm 68), USFWS Little White Salmon National Fish Hatchery
<i>Incubation location (facility name, stream, Rkm, subbasin)</i>	Klickitat River Hatchery (Rkm 68), USFWS Little White Salmon National Fish Hatchery
<i>Rearing location (facility name, stream, Rkm, subbasin)</i>	Klickitat River Hatchery (Rkm 68), and Wahkiacus Acclimation Facility (Rkm 27)

1.6 Type of program.

Segregated Harvest

No conservation goal has been established for Klickitat fall Chinook because this race of Chinook is not native to the Subbasin. There are no plans to establish a viable, naturally reproducing population of fall Chinook in the Subbasin in the foreseeable future. The goal is to establish a locally adapted, segregated hatchery population designed to provide fish for harvest. The program will support Tribal and non-tribal fisheries mandated by federal court orders and treaties. The objective is production of 18,000 fall Chinook for harvest in all fisheries, with the majority occurring in Tribal fisheries in Zone 6 and the Klickitat River.

The YN considered five options for managing Klickitat River fall Chinook.

1. Maintain existing program
2. Transition to fully integrated hatchery program
3. Eliminate hatchery production
4. Restore the natural fall Chinook spawning habitat eliminated by the construction of The Dalles and John Day dams
5. Convert existing program to local broodstock for ½ (2 million) on-station smolt release and partner with USFWS for broodstock collection within the Klickitat Subbasin (if needed) for pre-smolt transfer for the other ½ (2 million) of the production from the USFWS's Little White Salmon NFH to the Wahkiacus Acclimation Facility for release.

The hatchery strategy involves implementation of a segregated harvest program that uses a portion of the fall Chinook returning to the Klickitat River as its broodstock. This will be accomplished through: 1) elimination of eyed-egg transfers from Little White Salmon NFH, 2) development of locally adapted

broodstock for ½ (2 million) of the production 3) construction of the Wahkiacus Acclimation Facility (WAF), to received ½ (2 million) of the production from LWSNFH as pre-smolts for acclimation prior to release, 4) marking all juvenile fall Chinook, and 5) releasing a total of 4 million fall Chinook subyearlings at 50 to 80 fish per pound annually.

The current 4 million production volume has been essentially unchanged since 1986. It can be summarized as:

1986-2007 eyed eggs (4 million) transferred from Priest Rapids Hatchery to the Klickitat Hatchery (KH); 2008-to-present eye eggs transferred from the Little White National Fish Hatchery (LWSNFH) in the form of 3.5 million green eggs/1.5 million eyed supplied in October – December. Eyed eggs from LWSNFH are shipped into the KH in December.

Eggs are incubated and hatched on Klickitat spring water from late October to February, when they are ready to be ponded into (spring water supplied) Raceway banks A & B (10 ponds total, RW's 1-7 and 12-14) where they were reared to for volitional release at 75-80/ffp generally in the month of June.

Beginning in 2013 the current LWSNFH brood source is currently under transition to a 100% green egg transfer program where 5 million green eggs will be delivered to the KH to meet the 4.0M production goal.

1.7 Purpose (Goal) of program.

The goal is to establish a locally adapted, and regionally adapted, segregated hatchery population designed to provide fish for harvest. The harvest will support Tribal and non-tribal fisheries mandated by federal court orders and treaties. The objective is production of 18,000 fall Chinook for harvest in all fisheries, with the majority occurring in Tribal fisheries in Klickitat River and Zone 6.

1.8 Justification for the program.

The program will be operated to provide fish for tribal and sport harvest while minimizing adverse affects on listed fish (steelhead and bull trout). Fall Chinook releases in the Klickitat River have long been a part of *U.S. v. Oregon* management agreements (see 2008-2017 *U.S. v. Oregon* Management Agreement) and are part of efforts to move Mitchell Act production to upriver areas where production losses occurred.

1.9 List of program "Performance Standards".

See section 1.10 below.

1.10 List of program "Performance Indicators", designated by "benefits" and "risks".

1.10.1 Benefits:

Benefits		
Performance Standard	Performance Indicator	Monitoring & Evaluation
Provide Fish to Meet Columbia River fish Mgt. Plan (<i>US v Oregon</i>), production and harvest objectives.	Contribute to a meaningful harvest for sport, tribal and commercial fisheries. Objective is to provide 18,000 adult fish for harvest.	Contribution to fisheries will be estimated for each brood year released. Work with co-managers to manage adult fish returning in excess of broodstock needs
Smolt-to-adult survival (SAR)	SAR Value of 1%	SAR will be determined by counting tagged fish recovered at traps, broodstock collection facilities, sport and tribal fisheries and on the spawning grounds.
Straying of Klickitat River origin fish to other subbasins	Stray rate of less than 5%	Regional M&E efforts will be used to track the number and capture location of Klickitat River origin fish
Use of local origin fall Chinook for broodstock	For the on-station 2.0 million release program, broodstock will consist of 95% Klickitat origin fall Chinook. For the 2.0M subyearling release from WAF, broodstock will consist of returns to the USFWS LWNFH	DNA sampling will be utilized to ensure in-basin Chinook are captured for this program.
Maintain outreach to enhance public understanding, participation and support of YKFP hatchery programs	Provide information about agency programs to internal and external audiences. For example, local schools and special interest groups tour the facility to better understand hatchery operations. Off- station efforts may include festivals, classroom participation, stream adoptions and fairs.	Evaluate use and/or exposure of program materials and exhibits as they help support goals of the information and education program. Record on-station organized education and outreach events.
Program contributes to fulfilling tribal trust responsibility mandates and treaty rights	Follow pertinent laws, agreements, policies and executive and judicial orders on consultation and coordination with Native American tribal governments.	Participate in annual coordination meetings between the co-managers to identify and report on issues of interest, coordinate management, and review programs (FBD process).
Implement measures for broodstock management to maintain integrity and genetic diversity	Between 1,250 - 2,500 adults are collected throughout the spawning run in proportion to timing, age and sex composition of return. 1,250 for in-basin segregated program, and additional brood for transfer to LWSNFH to support (if needed) brood for per-smolt transfer.	Annual run timing, age and sex composition and return timing data are collected.
Region-wide, groups are marked in a manner consistent with information needs and protocols to estimate impacts to natural and hatchery-origin fish	16.5% of all fish released will be ad-clipped and coded-wire-tagged (cwt). Blank coded-wire tags may be used to identify fish for broodstock. 100% will be adipose-clipped.	Returning fish are sampled throughout their return for length, sex, and mark
Maximize survival at all life stages using disease control and disease prevention techniques. Prevent introduction, spread or amplification of fish pathogens. Follow Co-managers Fish Health Disease Policy (WDFW and NWIFC 1998).	a. Necropsies of fish to assess health, nutritional status, and culture conditions	USFWS Fish Health Pathologist inspect adult broodstock yearly for pathogens at Little White Salmon NFH and Klickitat Hatcheries and monitor juvenile fish on a monthly basis to assess health and detect potential disease problems. As necessary, USFWS Fish Health Pathologist recommends remedial or preventative measures to prevent or treat disease, with administration of therapeutic and prophylactic treatments as necessary

		A fish health database will be maintained to identify trends in fish health and disease and implement fish health management plans based on findings.
	b. Release and/or transfer exams for pathogens and parasites.	1 to 6 weeks prior to transfer or release, fish are examined in accordance with the Co-managers 'Fish Health Policy
	c. Inspection of adult broodstock for pathogens and parasites.	At spawning, lots of 60 adult broodstock are examined for pathogens
	d. Inspection of off-station fish/eggs prior to transfer to hatchery for pathogens and parasites.	Controls of specific fish pathogens through eggs/fish movements are conducted in accordance to Co-managers Fish Health Disease Policy (WDFW and NWIFC 1998)

1.10.2 Risks:

Risks		
Performance Standard	Performance Indicator	Monitoring & Evaluation
Minimize impacts and/or interactions to ESA listed fish	Hatchery operations comply with all state and federal regulations. Hatchery juveniles are raised to smolt-size (50-80 fish/lb) and released voluntarily from the hatchery and acclimation sites at a time that fosters rapid migration downstream. Also, 16.5 % of all fish released will be marked and CWT tagged to identify them from naturally produced fish and monitor straying. All fish will be adipose-clipped.	Monitor size, number, date of release and CWT mark quality.
Artificial production facilities are operated in compliance with all applicable fish health guidelines, facility operation standards and protocols including IHOT, Co-managers' Fish Health Policy and drug usage mandates from the Federal Food and Drug Administration	Hatchery goal is to prevent the introduction, amplification or spread of fish pathogens that might negatively affect the health of both hatchery and naturally reproducing stocks and to produce healthy smolts that will contribute to the goals of this facility.	Pathologists from USFWS Fish Health Section monitor program monthly. Exams performed at each life stage may include tests for virus, bacteria, parasites and/or pathological changes, as needed
Ensure hatchery operations comply with state and federal water quality and quantity standards through proper environmental monitoring	NPDES permit compliance YN water right permit compliance	Flow and discharge reported in monthly NPDES reports.
Water withdrawals and in-stream water diversion structures for hatchery facility will not affect spawning behavior of natural populations or impact juveniles.	Hatchery intake structures meet state and federal guidelines where located in fish bearing streams.	Barrier and intake structure compliance assessed and needed fixes are prioritized.
Hatchery operations comply with ESA responsibilities	YN completes an HGMP and is issued a federal and state permit when applicable.	
Harvest of hatchery-produced fish minimizes impact to wild populations	Harvest is regulated to meet appropriate biological assessment criteria. Mass mark juvenile hatchery fish prior to release to enable state agencies to implement selective fisheries.	Harvests are monitored by agencies and tribes to provide up to date information.

1.11.1 Proposed annual broodstock collection level (maximum number of adult fish).

The program will collect between 1,250 - 2,500 adults returning to the Klickitat River (1:1 Male to Female) for the local on-station program.

1.11.2 Proposed annual fish release levels (maximum number) by life stage and location.

Age Class	Max.	Size	Release	Location
-----------	------	------	---------	----------

	No. (Million)	(fpp)	Date	Stream	Release Point (RKm)	Major Water- shed	Eco- province
Fingerling	2.0	50.0 - 80.0	June/July	Klickitat	RKm 27	Klickitat	Columbia Gorge
Fingerling	2.0	50.0 - 80.0	June/July	Klickitat	RKm 68	Klickitat	Columbia Gorge

1.12 Current program performance, including estimated smolt-to-adult survival rates, adult production levels, and escapement levels. Indicate the source of these data.

Recent conversion from Priest Rapids Hatchery to Little White Salmon NFH (2007) do not present enough years of information so the following information from historic Priest Rapids egg source is presented. This information will be updated for this section in the future.

Smolt-to-adult survival rates for URB from Priest Rapids Hatchery fall Chinook have been estimated to range from 0.29 % to 2.44 % (smolt-to-adult overall survival estimates for brood years 1983-87 from IHOT 1995). In comparison, from 1995 – 1999, SARs for Klickitat URB have averaged 0.34% and ranged from a low of .009% in 1995 to a high 0.73% in 1999 (RMIS Database).

Smolt-to-adult survival rates as estimated by University of Washington Columbia Basin Research office (available at <http://www.cbr.washington.edu/trends/index.php>) for URB fall Chinook for brood years 2005-2008 averaged 0.24% with a range of 0.05% to 0.42%. Some years have incomplete recovery data so these estimates should be considered preliminary.

Converting ½ (2 million) of the total production to local broodstock is expected to result in an average SAR of 1% for that component which is closer to that observed for Priest Rapids Hatchery. This hatchery uses local origin fall Chinook as its brood source.

The average annual harvest of fall Chinook from Klickitat River releases in combined ocean, Columbia River, and Klickitat River fisheries is estimated to exceed 18,000 fish. Sport and Tribal fall Chinook fisheries in the Klickitat River take, on average, about 3,700 fish each year.

Harvest rates on aggregate Upper River Bright fall Chinook stocks (including Klickitat River fall Chinook) in ocean and lower Columbia River fisheries range from 40% to 60% (NOAA 2000). Klickitat River terminal harvest rates of URB fall Chinook averaged approximately 45% from 1986 to 2008.

1.13 Date program started (years in operation), or is expected to start.

The Klickitat Hatchery was completed in 1951 and fall Chinook production efforts have been on-going since that time. Beginning in 1986, Klickitat Hatchery

production switched from the earlier tule stock to an upriver bright (URB) fall Chinook from Priest Rapids. In 2007, converted URB egg source from Priest Rapids Hatchery to Little White Salmon NFH.

1.14 Expected duration of program.

On-going program. A Memorandum of Understanding (MOU) was signed on December 30, 2005 detailing the transfer of ownership and operational responsibility of the Klickitat Hatchery and the Lyle Falls and Castile Falls fishways from the WDFW to the YN. Overall goal is to maintain the Klickitat fall Chinook program at current levels for harvest augmentation for the future

1.15 Watersheds targeted by program.

The Klickitat River Subbasin and the mainstem Columbia River (harvest)

1.16 Indicate alternative actions considered for attaining program goals, and reasons why those actions are not being addressed.

A list of the alternative approaches considered for the program is presented below. More detailed discussions to why each alternative was rejected can be found in the revised Klickitat River Anadromous Fisheries Master Plan (Yakama Nation 2012 Draft).

Alternative 1- Maintain Existing Program: Eliminated due to: 1) imports fish from outside of the basin which increases disease risk, and 2) competition effects on spring Chinook.

Alternative 2- Transition to Fully Integrated Hatchery Program: Program requires the establishment of naturally reproducing population above Lyle Falls. Since fall Chinook were not historically present in this area, their presence may result in negative impacts to native spring Chinook.

Alternative 3- Eliminate Hatchery Production: Alternative did not meet the harvest goals identified in the *U.S. v. Oregon* 2008-17 Columbia River Fishery Management Plan. .

Alternative 4- Restore the Natural Fall Chinook Spawning Habitat Eliminated by the Construction of The Dalles and John Day Dams: Primarily eliminated due to political infeasibility.

Alternative 5- Convert existing program to local broodstock for ½ (2 million) of the production and partner with USFWS for pre-smolt transfer for the other ½ (2 million) of the production as in this HGMP.

1.16.3) Potential Reforms and Investments

Reform/Investment 1: Upgrade of adult collection facilities at Lyle Falls.

Reform/Investment 2: Construction of the Wahkiacus Acclimation Facility.

Reform/Investment 3: Addition of staff, equipment, and supplies needed to

implement program.

Reform/Investment 4: Release of $\frac{1}{2}$ (2 million) of total fall Chinook production lower in the Subbasin to reduce competition with listed fish species.

Reform/Investment 5: Use Klickitat River-origin adult returns as portion of the broodstock.

Section 2: Program Effects on ESA-Listed Salmonid Populations

2.1 List all ESA permits or authorizations in hand for the hatchery program.

Program is described in the Biological Assessment For The Operation Of Hatcheries Funded by The National Marine Fisheries Service (March 1999), Statewide Section 6 consultation with USFWS for interactions with Bull Trout, and concurrent with this HGMP, to satisfy Section 7 consultations the YN is writing HGMPs to cover all stock/programs in the Klickitat River including fall Chinook, spring Chinook, steelhead, and coho released from Klickitat Hatchery.

This document is intended to be consistent with NOAA (2008) which states (RPA 39):

The FCRPS Action Agencies will continue funding hatcheries in accordance with existing programs... Consultation under the ESA on the operation of hatchery programs funded by the FCRPS Action Agencies [will] include the submittal of updated and complete HGMPs. Updated and complete HGMPs are to be submitted to NOAA Fisheries and ESA consultation should be initiated by ... July 2009 for hatchery programs in the Middle Columbia ... ESA consultations should be completed by January 2010 for hatchery programs in the Middle Columbia ...

Project sponsors are also aware of direction in NOAA (2009) calling “for consultations on hatchery programs within the MCR Steelhead DPS to be completed by January 2010”. Project sponsors remind NOAA of its statement in this document that “mitigation obligations will not be diminished under this process”. The Yakama Nation considers this project essential to meeting federal commitments to honor the Treaty of 1855, and to “protect, rebuild, and enhance” anadromous salmon populations throughout tribal usual and accustomed fishing areas as described in the 2008-2017 United States v. Oregon Management Agreement and in the Columbia River Fish Accords. As such, any changes to program parameters which would diminish the number of adult salmon returning to tribal usual and accustomed fishing areas that result from this HGMP development and consultation process will not be implemented unless and until they are considered and approved in appropriate policy fora.

NOAA. 2008. Consultation Title: Remand of 2004 Biological Opinion on the Federal Columbia River Power System (FCRPS) including 19 Bureau of Reclamation Projects in the Columbia Basin (Revised pursuant to court order, NWF v. NMFS, Civ. No. CV 01-640-RE (D. Oregon). Tracking Number: 2005/05883. https://pcts.nmfs.noaa.gov/pls/pcts-pub/pcts_upload.summary_list_biop?p_id=27149


NOAA. 2009. Letter from Rob Jones, Chief, Salmon Recovery Division, National Marine Fisheries Service, Portland Oregon to “Interested Parties”, dated March 5, 2009. NMFS, Portland Office, 1201 NE Lloyd Blvd, Suite 100, Portland, Oregon.

2.2 Descriptions, status and projected take actions and levels for ESA-listed natural populations in the target area.

ESA listed stock	Status	Take Level	Action
Summer Steelhead-Natural	Threatened	Minor	Broodstock collection/trapping at Lyle Falls
Winter Steelhead-Natural	Threatened	Minor	Broodstock collection/trapping at Lyle Falls
Bull Trout – Natural	Threatened	Minor	Broodstock collection/trapping at Lyle Falls

2.2.1 Description of ESA-listed salmonid population(s) affected by the program.

Adult and juvenile run-timing for listed steelhead and other fish species are presented in the figure below.



The majority of the steelhead population is found from the mouth of the Klickitat River to Castile Falls. Until improvements in 2005, steelhead access to areas above Castile falls had been limited due to poor migration conditions at the Castile Falls fishway tunnels. Steelhead spawning is concentrated between Rkm 8 and 80. Tributary spawning occurs in White Creek watershed (including Brush and Tepee Creeks), Summit Creek, Dead Canyon Creek, the lower Little Klickitat watershed (including Bowman and Canyon Creeks), Swale Creek, Snyder Creek, and occasional use of tributaries below the town of Klickitat.

Juvenile rearing occurs in the mainstem and major tributaries. Peak smolt migration occurs in April and May; however, juvenile steelhead have been captured in traps located at Lyle Falls in all months.

Klickitat River bull trout life history characteristics are not very well understood, but bull trout are present in the middle and lower mainstem Klickitat and are concentrated in the West Fork Klickitat watershed and several of its tributaries. The population in the West Fork Klickitat watershed is likely resident and while the fish found in the mainstem Klickitat are likely adfluvial or migratory. Falls on the West Fork likely isolate most of the resident bull trout population from the mainstem Klickitat River.

Maps depicting steelhead and bull trout distribution in the Klickitat River are presented in Appendix 2.

Identify the ESA-listed population(s) that will be directly affected by the program

No NMFS ESA listed fish populations will be directly affected by this program. This broodstock was not considered part of the ESU by WDFW and USFWS and was not essential for recovery. This stock originates from populations not considered to be part of the Lower Columbia River Chinook salmon ESU.

Identify the ESA-listed population(s) that may be incidentally affected by the program

Middle Columbia River Steelhead March 19, 1998; 64 FR 14508; updated January 5, 2006, 71 FR 834. Threatened
Columbia Basin DPS Bull Trout June 10, 1998 (63 FR 31647), Threatened.

2.2.2. Status of the ESA-listed population(s) affected by the program

Middle Columbia River Steelhead (*Oncorhynchus mykiss*) March 19, 1998; 64 FR 14508; updated January 5, 2006, 71 FR 834 Threatened.

The ICTRT (2007) has identified Klickitat River steelhead as an independent population belonging to the Mid-Columbia ESU. The Middle Columbia steelhead ESU was listed as threatened under the ESA on March 25, 1999 (64 FR 14517) and reaffirmed January 5, 2006; (71 FR 834). The Klickitat steelhead population includes both summer-run and winter-run steelhead (Yakima Nation 2012).

Temporal and spatial spawning segregation between the Klickitat steelhead summer and winter runs has not yet been clearly defined. Ongoing genetic analysis and radio telemetry monitoring is expected to provide additional information about the spatial and temporal distribution of both steelhead races.

Past genetic analysis on steelhead have shown some degree of genetic differentiation between tributaries to the Klickitat River; genetic samples from the upper Klickitat, White Creek, and Trout Creek seem to diverge most widely from the Skamania Hatchery stock (Marshall 2000). Recent genetic analysis indicates there may be six to seven genetically distinct populations of naturally reproducing steelhead in this river system. The results also suggest the genetic integrity and

variation of native Klickitat River steelhead have been maintained despite repeated hatchery introduction and that the potential is high for restoring the population's viability (Narum et al. 2006).

No solid historical data exist on the size and productivity of the Klickitat summer steelhead run. Based on NOAA Fisheries historical intrinsic potential analysis, the ICTRT considers the Klickitat River population to be an "intermediate" sized population that can support a minimum of 1,000 spawners (ICTRT 2007).

The escapement of naturally spawning (summer and winter, hatchery and wild combined) steelhead in the Klickitat River from 1987 to present has been estimated at approximately 700 fish (see below). However, this estimate is based on redd count data which is believed to be an underestimate because of difficulties associated with conducting accurate counts during spring flow conditions (NPCC 2004). YN biologists hypothesize that the actual mean escapement is closer to 900-1000 spawners annually.

Recent mark-recapture evaluations using hatchery and natural origin summer steelhead trapped and tagged at Lyle Falls estimates natural origin returns to the lower Klickitat River to average about 1600 fish from 2005-2011 (Gray 2007 and Zendt et al. 2013).

Additionally, from the early 1960s to 2005, Castile Falls likely blocked all steelhead from stream habitat located upstream of the falls. By 2005, upstream fish passage conditions at Castile Falls were improved to allow steelhead access to this portion of the Subbasin. Habitat modeling work indicates that adult steelhead production potential above the falls may be as high as 750 adults (Yakama Nation 2012). If the production potential estimate is accurate, total steelhead production in the Klickitat River Subbasin could increase to over 2,000 fish in the near future.

Based on population parameters developed for the area mainly below Castile Falls, the ICTRT rated Klickitat steelhead as having only a moderate risk in regards to the key population parameters of abundance/productivity and spatial structure/diversity. Thus, the population does not meet ICTRT criteria for a viable population, although it does meet criteria for a "Maintained" population (ICTRT 2007).

Columbia Basin DPS Bull Trout (*Salvelinus confluentus*) June 10, 1998 (63 FR 31647), Threatened.

The Fish and Wildlife Service issued a final rule listing the Columbia River and Klamath River populations of bull trout (*Salvelinus confluentus*) as a threatened species under the Endangered Species Act on June 10, 1998 (63 FR 31647). The Columbia River Distinct Population Segment is threatened by habitat degradation and fragmentation, blockage of migratory corridors, poor water quality, and past fisheries management practices such as the introduction of non-native species.

The Lower Columbia Recovery Unit Team identified two core areas (Lewis and Klickitat rivers) within the recovery unit. The Klickitat Core Area includes all tributaries downstream to the confluence with the Columbia River. Recent

evidence indicates both resident and adfluvial bull trout are present in the Subbasin. Numerous confirmed and anecdotal reports of bull trout exist in the mainstem Klickitat River from the mouth up to the area below Castile Falls. Sizes reported are indicative of an adfluvial life history. Presence of resident populations has also been documented in the West Fork Klickitat River, Fish Lake Stream, Little Muddy Creek, Trappers Creek, Clearwater Creek, Two Lakes Stream, and an unnamed tributary to Fish Lake Stream (all within the West Fork Klickitat watershed) (Byrne et al. 2001, Thiesfeld et al. 2002, Gray 2007).

The abundance of the stock in the Klickitat River is poorly known and there are insufficient data to make an assessment. However, it appears that there are very few bull trout in the lower- to mid-Klickitat drainage. Bull trout appear to be more abundant in the upper drainage where habitat conditions are more favorable.

Preliminary results of recent genetic analysis indicate that resident bull trout in the Klickitat Subbasin are genetically distinct from other Columbia tributary populations, but that fish in two West Fork Klickitat tributaries (Trappers and Clearwater creeks) do not differ significantly from each other.

The impacts of hatchery salmon and steelhead in the main Klickitat River on bull trout are unknown. Generally, in drainages colonized by anadromous salmon and steelhead, char successfully co-exist by occupying a different ecological niche. However, negative interactions such as predation may occur when hatchery fish are released near char spawning and rearing areas.

2.2.3 Describe hatchery activities, including associated monitoring and evaluation and research programs, that may lead to the take of listed fish in the target area, and provide estimated annual levels of take.

The following activities may lead to a take of listed species.

Broodstock Collection: Broodstock will be collected for this program at Lyle Falls and Klickitat Hatchery from September through November. No ESA-listed fish mortalities (steelhead or bull trout) have been observed at the Klickitat River Hatchery for the past nine years. The operation of the new adult collection facilities at Lyle Falls may result in some ESA- listed steelhead being handled because the collection facilities will be operated during time of the year when steelhead may be migrating to spawning grounds. These two facilities have been constructed to meet NMFS passage and handling criteria which will minimize stress and associated mortality rates on fish.

Water diversion: Water is diverted at times from the Klickitat River for Klickitat Hatchery operations. This diversion affects approximately 0.25 miles of stream habitat. The loss in habitat may result in a decrease in steelhead and bull trout abundance, although this has not been quantified, it is expected to be negligible. In addition, NOAA (2008) (http://www.nwr.noaa.gov/Salmon-Hydropower/FERC/upload/Fish_passage_design.pdf) has identified design and alternatives needed to bring existing structures in compliance with NOAA fish screening standards. YN has requested funding for future scoping, design, and

construction work of a new intake system.

The new WAF will divert approximately 24 cfs of water from the Klickitat River for rearing both fall Chinook and coho. The diversion structure will be screened to meet NMFS standards. The 24 cfs equates to approximately 1% of total river flow in May, the peak usage month. The diverted water will decrease streamflow in less than 0.10 miles of riverine habitat.

Water Quality: Both the Klickitat Hatchery and AF will operate under the “*Federal Aquaculture Facilities and Aquaculture Facilities Located in Indian Country within the Boundaries of the State of Washington*” National Pollution Discharge Elimination System (NPDES) general permit (WAG-130021) which conducts effluent monitoring and reporting and operates within the limitations established in its permit administered by the U.S. Environmental Protection Agency (EPA). Monthly and annual reports on water quality sampling, use of chemicals at this facility, compliance records are available from EPA. Discharges from the cleaning treatment system are monitored as follows:

Total Suspended Solids (TSS): Collected 1 to 2 times per month on composite effluent, maximum effluent, and influent samples.

Settleable Solids (SS): Collected 1 to 2 times per week on effluent and influent samples.

In-hatchery Water Temperature - Daily maximum and minimum readings. Water quality monitoring is not expected to result in the take of listed species.

Genetic introgression: Straying of fall Chinook from this program to other subbasins could result in genetic introgression with listed Chinook stocks, e.g. Snake River fall Chinook. Indirect take from straying is unknown.

Disease: Outbreaks in the hatchery may cause significant adult, egg, or juvenile mortality. Over the years, advances in rearing densities, disease prevention, and fish health monitoring have greatly improved the health of the programs at Klickitat River Hatchery. Policies and Procedures for Columbia Basin Anadromous Salmonid Hatcheries (IHOT 1995) have been instrumental in reducing disease outbreaks. Fish are planted and transferred after a fish health specialist has determined population health. Indirect take from disease is unknown.

Release Program

Competition and Predation: According to the HSRG (2005) and Flagg et al. (2000) the potential for predation of wild salmonids by hatchery-reared smolts will depend on the size, number, and spatial distribution of both predators and prey, the functional and numerical responses of the predators, and the amount of time that predators and prey are in proximity. Busack et al. (2005) reviewed published rates of predation by juvenile hatchery fall Chinook on wild juvenile Chinook and found predation rates were generally low (<2% of natural population consumed). In contrast, data collected on hatchery coho predation rates on wild fall Chinook

juveniles in the Lewis River were quite high (>11%) (Hawkins and Tipping 1999). The variability in study results is one reason the HSRG (2005) suggests that hatcheries monitor predation impacts resulting from hatchery releases.

In general, hatchery fish can consume fish that are 50% of their body length, however studies reviewed by Busack et al. (2005) indicated that the range may extend from approximately 38% (steelhead) to 75% (coho). NOAA Fisheries and the USFWS in a number of biological assessments and opinions were of the opinion that juvenile salmonids can consume prey up to 33% of their body length (USFWS 1994; NMFS 1999). Predation by hatchery fish on wild fish can occur anywhere the two stocks exist in space and time. Therefore, predation may not only be a concern in the stream environment, but also in the estuary and marine environments.

The site-specific nature of predation and the limited number of empirical studies that have been conducted make it difficult to predict the predation effects of this specific hatchery release. The YN is unaware of any studies that have empirically estimated the predation risks to listed fish posed by the Klickitat Hatchery programs. In the absence of site-specific empirical information, the identification of risk factors can be a useful tool for reviewing hatchery programs while monitoring and research programs are developed and implemented in the region.

Risk Factors:

Date of Release: The release date can influence the likelihood that listed species are encountered. Fall Chinook will be released in June, which is within the window when steelhead fry will be present in the mainstem Klickitat River.

Fish Size at Release: Based on the 33% of body length predation assumption and a fall Chinook size of release range of 80-100 mm, hatchery Chinook may consume listed steelhead up to 26-33 mm in length. During the time fall Chinook are expected to be in the mainstem Klickitat River (June-July), steelhead fry (26-40 mm) will be present in the system and will be vulnerable to predation. However, steelhead fry will be occupying shallow water habitat that is not likely accessible to the larger hatchery fall Chinook; thereby reducing the chance of predation. The level of fall Chinook predation on steelhead is unknown.

Release Location and Release Type: The likelihood of predation may also be affected by the location and the type of release. Other factors being equal, the risk of predation may increase with the length of time that fish co-mingle. In the freshwater environment, this is likely to be affected by distribution of the listed species in the watershed, the location of the release and the speed at which fish released from the program migrate. Fall Chinook will be released volitionally from rearing sites located at Rkm 27 (WAF). The remaining Fall Chinook production (2,000,000) will be released volitionally from the Klickitat Hatchery.

The minimal amount of time they spend in the river system should reduce predation and competition effects to listed fish species. Additionally, as fall Chinook will not be released in tributaries, they will not affect steelhead juveniles rearing in these streams.

Residualism: For the fall Chinook subyearling program the following actions are

taken to reduce residualism:

1. Fish Condition factors, standard deviation and co-efficient of variation (CV) on lengths of fish are measured throughout the rearing cycle and at release.
2. Feeding rates and regimes throughout the rearing cycle are programmed to satiation feeding to minimize size variations and reprogrammed as needed to achieve goals for smolt size at time of release.
3. Based on past history, fish have reached a size and condition that indicates a smolted condition at release.
4. Releases occur within known time periods of wild fish migration.
5. Releases from acclimation ponds are volitional with large proportions of the populations moving out initially with the remainder of the population vacating within a couple of days.

Straying: As noted by the HSRG (2009), straying to the Snake River of fall Chinook released in the Klickitat has been a major concern in the past. However, past releases in the Klickitat used Priest Rapids (and in some years even Lyons Ferry Hatchery) as brood sources. Beginning in brood year 2007, the URB brood source is from the Little White Salmon River and approximately 650,000 fish or 16.25% of the Klickitat release is coded-wire tagged (per US v. Oregon). This has helped better determine minimal stray rates of Klickitat releases of fall Chinook return upstream to Ice Harbor Dam on the Snake River. Through production changes outlined in this HGMP further attempts are underway to address this issue by developing a locally adapted broodstock and releasing 50% of the fish (~2.0 million) from an acclimation site at Wahkiacus. Marking methods will allow continued monitoring of this issue.

Migration Corridor/Ocean:

The Columbia River hatchery production ceiling, called for in the Proposed Recovery Plan for Snake River Salmon of approximately 197.4 million fish (1994 release levels), has been incorporated by NOAA-Fisheries into their recent hatchery biological opinions to address potential mainstem corridor and ocean effects, as well as other potential ecological effects from hatchery fish. Recent releases have been in the 140 to 145 million range for the entire Columbia River basin. Although hatchery releases occur throughout the year, approximately 80% occur from April to June and Columbia River mainstem out-migration occurs primarily from April through August (www.fpc.org). It is unknown to what extent listed fish are available both behaviorally and spatially on the migration corridor. Once in the main stem Columbia River, Witty et al. (1995) has concluded that predation by hatchery fish on wild salmonids does not significantly impact naturally produced fish survival in the Columbia River migration corridor. In a study designed to define the migrational characteristics of Chinook salmon, coho salmon, and steelhead trout in the Columbia River estuary, Dawley et al (1984),

found the average migration rates for subyearling Chinook, yearling Chinook, and coho salmon and steelhead, were 22, 18, 17, and 35 Rkm/d respectively. There appear to be no studies demonstrating that large numbers of Columbia system smolts emigrating to the ocean affect the survival rates of juveniles in the ocean in part because of the dynamics of fish rearing conditions in the ocean and an inability to measure.

Monitoring:

Monitoring and evaluation activities have the potential to harass, kill or injured handled fish; these activities are summarized below and described in more detail in Appendix 1.

Monitoring and evaluation activities include: spawning ground surveys; adult salmonid monitoring at Lyle Falls and Castile Falls fishways; juvenile outmigration monitoring (using floating rotary screw traps); juvenile and resident salmonid population surveys (via stream electrofishing); scale analysis; sediment monitoring; temperature and water quality monitoring; habitat surveys; and genetic data collection and analysis.

Research:

No research program is associated with the fall Chinook hatchery program, other than the monitoring and evaluation activities described above and in Appendix 1.

Provide projected annual take levels for listed fish by life stage (juvenile and adult) quantified (to the extent feasible) by the type of take resulting from the hatchery program (e.g. capture, handling, tagging, injury, or lethal take).

Data on the take of listed species is presented in the following table. These numbers include both natural- and hatchery-origin steelhead found in the Klickitat River subbasin. Hatchery-origin steelhead released in and returning to the Klickitat River are primarily from Skamania Hatchery. A breakdown of estimated take associated with each M&E activity is presented in Appendix 1.

Estimated listed salmonid take levels by hatchery activity.

Steelhead

<i>ESU/Population</i>	Middle Columbia River Steelhead
<i>Activity</i>	Klickitat Hatchery Spring Chinook Program
<i>Location of hatchery activity</i>	Klickitat River Hatchery
<i>Dates of activity</i>	Year-round
<i>Hatchery Program Operator</i>	YN & WDFW through co-managed YKFP

Type of Take	Annual Take of Listed Fish by life Stage (number of fish)
--------------	--

	Egg/Fry	Juvenile /Smolt	Adult	Carcass
Observe or harass (a)		50-150	100	
Collect for transport (b)				
Capture, handle, and release (c)		500-2000 wild; 2000-3000 hatchery*	650-950 wild; 800-1000 hatchery**	
Capture, handle, tag/mark/tissue sample, and release (d)		3000-10,000 wild***		
Removal (e.g., broodstock) (e)				
Intentional lethal take (f)				
Unintentional lethal take (g)		190 wild; 150 hatchery	25 wild; 30 hatchery	
Other take (indirect, unintentional) (h)				

* Smolt trapping operations for monitoring purposes

** Although steelhead have not been taken during past hatchery practices, it is anticipated that adult steelhead will be collected and handled at the new collection facilities at Lyle Falls. Mortality occurs only on rare occasions during these operations.

***Stream fish (juvenile steelhead and resident trout) sampling operations

- a. Contact with listed fish through stream surveys, carcass and mark recovery projects, or migrational delay at weirs.
- b. Take associated with weir or trapping operations where listed fish are captured and transported for release.
- c. Take associated with weir or trapping operations where listed fish are captured, handled and released upstream or downstream.
- d. Take occurring due to tagging and/or bio-sampling of fish collected through trapping operations prior to upstream or downstream release, or through carcass recovery programs.
- e. Listed fish removed from the wild and collected for use as broodstock.
- f. Intentional mortality of listed fish, usually as a result of spawning as broodstock.
- g. Unintentional mortality of listed fish, including loss of fish during transport or holding prior to spawning or prior to release into the wild, or, for integrated programs, mortalities during incubation and rearing.
- h. Other takes not identified above as a category.

Indicate contingency plans for addressing situations where take levels within a given year have exceeded, or are projected to exceed, take levels described in this plan for the program.

Any mortality from this operation or other Klickitat River Hatchery operations will be communicated to Fish program staff for additional guidance. For other listed species, if significant numbers of wild salmonids are observed to be negatively impacted by this operation, staff will inform the YN lead biologist who will communicate concerns to NOAA staff. Mitigation recommendations made by NOAA and the co-managers will then be implemented.

Section 3: Relationship of Program to Other Management Objectives

3.1 Describe alignment of the hatchery program with any ESU-wide hatchery plan (e.g. *Hood Canal Summer Chum Conservation Initiative*) or other regionally accepted policies (e.g. the *NPPC Annual Production Review Report and Recommendations - NPPC document 99-15*). Explain any proposed deviations from the plan or policies.

For ESU-wide hatchery plans, the plant of fall Chinook to the Klickitat River is consistent with:

- 1999 Biological Opinion on Artificial Propagation in the Columbia River Basin (NMFS 1999)
- Policies and Procedures for Columbia Basin Anadromous Salmonid Hatcheries (IHOT 1995)
- The 2008-2017 *U.S. v. Oregon* Management Agreement
- Columbia River Basin Fish and Wildlife Program (<http://www.nwcouncil.org/library/2000/2000-19/Default.htm>)
- NPPC Annual Production Review
- Principles and Recommendations of the HSRG (HSRG 2005 and 2009)
- Yakima/Klickitat Fisheries Project (YKFP or Project)
- Klickitat River Anadromous Fisheries Master Plan (2012, in draft)
- Recovery Plan for the Klickitat River Population of the Middle Columbia River Steelhead Distinct Population Segment (NOAA-Portland 2009).
- 2008 Columbia Basin Fish Accords Memorandum of Agreement between the Three Treaty Tribes and FCRPS Action Agencies

The current program has been made consistent with and aligned with the following plans and policies to the extent possible:

Yakima/Klickitat Fisheries Project (YKFP or Project) - Encompasses both the Yakima and Klickitat subbasins. It is the only major project in the Northwest Power and Planning Council's (NPPC) Fish and Wildlife Program that covers two major subbasins, each within a separate province. Since inception, the Yakama Nation has managed Project operations in both subbasins as one undertaking. By consolidating management for both subbasins into a single management unit, the YN has ensured Project efficiency at all levels. As necessitated by the NPPC's provincial proposal format, this proposal "unbundles" Project operation and maintenance activities. It covers the Klickitat Subbasin only. The YKFP is a supplementation project designated by the Northwest Power Planning Council's as the principal means of protecting, mitigating, and enhancing the anadromous fish populations in the Yakima and Klickitat subbasins (<http://www.ykfp.org/>).

Klickitat Master Plan -Prepared by Yakama Nation and reviewed by WDFW, this master plan addresses proposed facilities, production protocols, monitoring and evaluation, and habitat improvements needed to manage spring and fall Chinook salmon, coho, steelhead, bull trout, and Pacific Lamprey in the Klickitat

Subbasin.

NMFS recently adopted a recovery plan for the Mid-Columbia Steelhead Distinct Population Segment (DPS). The NMFS *Mid-Columbia Steelhead Recovery Plan (NMFS 2009)* summarizes information from four regional management unit plans covering the range of tributary habitats associated with the DPS in Washington and Oregon. Each of the management unit plans are incorporated as appendices to the recovery plan, along with modules for the mainstem Columbia hydropower system and the estuary, where conditions affect the survival of steelhead production from all of the tributary populations comprising the DPS. The recovery objectives defined in the plan are based on the biological viability criteria developed by the ICTRT. The plan also incorporates information on current status developed through the ICTRT (Ford *et al.* 2010).

U.S. v Oregon and the Columbia River Fish Management Plan (CRFMP)

Genetic Manual and Guidelines for Pacific Salmon Hatcheries in Washington. These guidelines define practices that promote maintenance of genetic variability in propagated salmon (Hershberger and Iwamoto 1981). Also, *Policies and Procedures for Columbia Basin Anadromous Salmonid Hatcheries* (Genetic Policy Chapter 5, IHOT 1995).

Spawning Guidelines for Washington Department of Fisheries Hatcheries. Assembled to complement the above genetics manual, these guidelines define spawning criteria to be used to maintain genetic variability within the hatchery populations (Seidel 1983). Also, *Policies and Procedures for Columbia Basin Anadromous Salmonid Hatcheries* (Genetic Policy Chapter 7, IHOT 1995).

Stock Transfer Guidelines. This document provides guidance in determining allowable stocks for release for each hatchery. It is designed to foster development of locally-adapted broodstock and to minimize changes in stock characteristics brought on by transfer of non-local salmonids (WDF 1991).

Fish Health Policy in the Columbia Basin. Details hatchery practices and operations designed to stop the introduction and/or spread of any diseases within the Columbia Basin.

National Pollutant Discharge Elimination System Permit Requirements (NPDES). This permit sets forth allowable discharge criteria for hatchery effluent and defines acceptable practices for hatchery operations to ensure that the quality of receiving waters and ecosystems associated with those waters are not impaired.

3.2 List all existing cooperative agreements, memoranda of understanding, memoranda of agreement, or other management plans or court orders under which program operates.

The program described in this HGMP is consistent with the following agreements and plans:

- The Columbia River Fish Management Plan
- Klickitat Master Plan

- Yakima/Klickitat Fisheries Project (YKFP or Project)
- U.S. vs. Oregon court decision and 2008-2017 Management Agreement
- Production Advisory Committee (PAC)
- Technical Advisory Committee (TAC)
- Integrated Hatchery Operations Team (IHOT) Operation Plan 1995 Volume III.
- Pacific Northwest Fish Health Protection Committee (PNFHPC) (<http://www.fws.gov/pnfhpc/>)
- In-River Agreements: State, Federal, and Tribal representatives
- Northwest Power Planning Council Sub Basin Plan (<http://www.nwcouncil.org/fw/subbasinplanning/klickitat/plan/>)
- Memorandum of Understanding Joint Operating Agreement for the Klickitat Hatchery (WDFW and YIN)
- 2008 Columbia Basin Fish Accords Memorandum of Agreement between the Three Treaty Tribes and FCRPS Action Agencies

3.3 Relationship to harvest objectives.

The *U.S. v. Oregon* Columbia River Fish Management Plan recognized the importance of tribal harvest in the Klickitat River by mandating releases of 4.0 million fall Chinook (<http://www.critfc.org/legal/crfmp88.html>). The YN has an overall objective of providing 18,000 adults for harvest. The fall Chinook program has provided a steady contribution to tribal commercial fisheries. A summary of fall Chinook escapement and harvest estimates are provided below in Section 3.3.1 for marine, Columbia River and Klickitat River. The data indicate that the fall Chinook program has provided, on average, approximately 18,600 fish for harvest.

3.3.1. Describe fisheries benefiting from the program, and indicate harvest levels and rates for program-origin fish for the last twelve years, if available.

The largest harvest of Klickitat Hatchery fall Chinook is in the Canadian Troll fishery, but they are also harvested in the Canadian sport and net fisheries, the Washington/Oregon coastal sport and troll fisheries, Alaskan fisheries, Columbia River tribal fisheries, and freshwater sport fisheries. Based on reported coded-wire tag recoveries through recovery year 2010, we estimate that harvest of fall Chinook produced or released from the Klickitat River annually averaged approximately 10,800 in marine fisheries, 6,000 in mainstem Columbia River fisheries (about 37% of the estimated average annual Columbia River mouth run size of Klickitat fall Chinook), and 3,800 in terminal Klickitat River fisheries (about 39% of the estimated average annual Klickitat River mouth run size of Klickitat fall Chinook) over the past 20-25 years. From 2000-2010, total harvest of Klickitat River fall Chinook in all fisheries combined ranged from about 11,500 to over 43,000 fish (and possibly higher in recent preliminary estimates) and was estimated to average approximately 28,000 fish annually. The tribal terminal fishery at Lyle Falls, one of the few remaining

traditional fishing sites for Yakama Nation fishers, harvested between 1,400 and 8,900 fall Chinook annually from 2000-2010.

Estimated harvest and run size information for Klickitat River fall Chinook (adults and jacks, early-fall/tule and URB stocks combined) (1986-2010).

Year	Marine ¹ Harvest	Columbia R. Mouth Return	Col. R. Harvest ²		Bonn. Passage Loss ³	Klickitat R. Mouth Return ⁴	Klickitat Harvest ⁴	Total Harvest
			Zones 1-5	Zone 6				
1986		25,702	9,020	7,451	461	8,768	8,106	24,578
1987		5,502	2,214	1,348	97	1,843	1,136	4,698
1988		3,705	1,694	878	57	1,076	1,076	3,648
1989	9,305	14,185	6,744	2,863	229	4,349	1,757	20,669
1990	15,689	12,817	4,109	2,952	288	5,468	1,574	24,324
1991	3,686	10,349	2,763	1,547	302	5,737	2,791	10,787
1992		7,687	1,694	1,169	241	4,583	1,148	4,011
1993	23,355	6,520	1,339	1,407	189	3,586	1,118	27,218
1994	1,015	6,686	296	731	283	5,377	1,249	3,290
1995	3,428	5,282	308	539	222	4,213	1,470	5,744
1996	4,482	13,922	1,679	1,541	535	10,166	3,809	11,512
1997	3,777	16,664	2,257	1,839	628	11,940	3,612	11,485
1998	3,201	18,007	2,293	1,433	714	13,566	3,504	10,432
1999	5,742	23,240	2,807	1,585	942	17,906	3,335	13,469
2000	4,383	21,372	3,205	2,980	759	14,428	4,939	15,507
2001	3,625	12,653	2,304	2,637	386	7,327	2,897	11,463
2002	19,337	33,609	5,881	7,194	1,027	19,507	7,742	40,154
2003	21,276	48,312	9,530	8,257	1,526	28,999	3,952	43,015
2004	25,217	22,979	3,278	3,628	804	15,269	9,086	41,209
2005	23,751	20,029	2,886	3,543	680	12,920	8,114	38,294
2006	11,371	14,929	806	3,646	524	9,954	6,201	22,024
2007	11,971	10,129	1,625	2,736	288	5,479	3,747	20,080
2008	10,353	13,733	2,458	3,772	375	7,129	3,965	20,547
2009	44,052	20,781	3,752	4,786	612	11,630	4,868	57,459
2010	49,702	24,365	2,365	4,487	876	16,637	11,471	68,025
Avg ⁵	10,788	16,001	3,095	2,855	502	9,547	3,753	18,616

1. Derived from Regional Mark Information System (RMIS) recovery year data for marine and freshwater coded-wire tag (CWT) recoveries of fall Chinook released in the Klickitat River.

2. Derived from *U.S. v. Oregon* Technical Advisory Committee reports.

3. Assume 5% passage attrition ascending Bonneville Dam and through the reservoir.

4. YN and WDFW database estimates.

5. Average excludes 2009-2010 (preliminary CWT estimates subject to change).

3.4 Relationship to habitat protection and recovery strategies.

The program described in this HGMP is consistent with the following habitat and protection strategies:

Klickitat Subbasin Recovery Plan for the Mid Columbia ESU- This plan provides habitat strategies to be used to recover ESA listed steelhead in the

Klickitat Subbasin. The hatchery program has considered current and future habitat conditions in sizing program and defining release locations.

Yakama Nation Fisheries Program (YNFP):

The Klickitat Watershed Enhancement Project is a BPA-funded watershed restoration project implemented by the Yakama Nation Fisheries Program (YNFP). The YNFP is working in coordination with WDFW, Natural Resources Conservation Service (NRCS), local Conservation Districts, local land trusts, and Regional Fisheries Enhancement Groups. The project was proposed under the Northwest Power Planning Council's Fish and Wildlife Program and funded by BPA in 1997. The project also solicits and receives significant funding from the Washington Salmon Recovery Funding Board. Initial project restoration projects were located within the Swale Creek and Little Klickitat River watersheds; ongoing projects focus on floodplain and riparian restoration in the mainstem Klickitat and the White Creek watershed. Included in the project scope of work are in-stream structural modifications, re-vegetation of the riparian corridor, floodplain reconnection, and exclusion fencing to prevent channel degradation from livestock. A monitoring program has been initiated to document project success and guide future restoration activities. Future phases of the project will use physical habitat survey and EDT modeling output to guide and prioritization restoration activities.

Klickitat River Anadromous Fisheries Master Plan:

The Master Plan is currently in preparation by the Yakama Nation in cooperation with WDFW. It addresses proposed goals and objectives, facilities, production, monitoring and evaluation, and habitat improvements needed to manage spring and fall Chinook salmon, coho salmon, steelhead, and Pacific Lamprey in the Klickitat Subbasin (Yakama Nation 2012 Step II Draft).

Subbasin Planning and Salmon Recovery:

A regional Subbasin planning process is a broad-scale initiative that will provide building blocks of recovery plans for listed fish. The fall Chinook hatchery program is designed to be consistent with the goals identified in this plan (NPPC 2004).

3.5 Ecological interactions.

Below are discussions on both negative and positive interactions relative to the fall Chinook program.

(1) Salmonid and non-salmonid fishes or species that could negatively impact the program:

Klickitat fall Chinook smolts are subject to predation after release throughout the entire migration corridor from the Subbasin to the mainstem Columbia River and estuary. Northern pikeminnows and introduced spiny rays as well as a variety of bird species including gulls, mergansers, cormorants, belted kingfishers, great blue herons and night herons are among those preying on fall Chinook smolts. River otters can take a heavy toll on migrating smolts while returning adults are preyed upon by harbor seals, sea lions, and Orcas. Large numbers of northern pikeminnows congregate at the mouth of the Klickitat River.

Predation on the juvenile Chinook outmigrants by the northern pikeminnow may have a negative impact on this stock. Avian predation by common mergansers, double crested cormorants, and (especially) Caspian terns also pose a large threat.

(2) Salmonid and non-salmonid fishes or species that could be negatively impacted by the program:

Natural salmon and steelhead populations that co-exist in local tributary areas and the Columbia River mainstem corridor areas could be negatively impacted by program fish. Of primary concern are the ESA-listed endangered and threatened salmonids: Snake River fall-run Chinook salmon ESU (threatened); Snake River spring/summer-run Chinook salmon ESU (threatened); Lower Columbia River Chinook salmon ESU (threatened); Upper Columbia River spring-run Chinook salmon ESU (endangered); Columbia River chum salmon ESU (threatened); Snake River sockeye salmon ESU (endangered); Upper Columbia River steelhead ESU (endangered); Snake River Basin steelhead ESU (threatened); Lower Columbia River steelhead ESU (threatened); Middle Columbia River steelhead ESU (threatened); and the Columbia River distinct population segment of bull trout (threatened). Listed fish can be impacted through a complex web of short- and long-term processes and over multiple time periods which makes evaluation of the net effect difficult. See previous discussion on potential ecological interactions in Section 2.2.3.

3) Salmonid and non-salmonid fishes or other species that could positively impact the program.

Other wild and hatchery salmonids may provide nutrients to the Klickitat River upon their return as adults. These carcasses may increase stream productivity, which in turn may increase food abundance for Chinook.

4) Salmonid and non-salmonid fishes or species that could be positively impacted by the program.

Aquatic and terrestrial species that consume salmonids will benefit from the continued release of fish from this program. Common species that may benefit include northern pikeminnow, smallmouth and largemouth bass, gulls, mergansers, cormorants, belted kingfishers, great blue herons, night herons, bald eagles, harbor seals, sea lions, river otters, bear, and killer whales (Orcas). Additionally, salmon carcasses act as a source of fertilizer that benefits riparian plants which, in turn, provide nutrients back to the stream.

Section 4. Water Source

4.1 Provide a quantitative and narrative description of the water source (spring, well, surface), water quality profile and natural limitations to production attributable to the water source.

Klickitat Hatchery

The Klickitat River is the primary homing water source for the target population. The water flowing into Klickitat Hatchery Pond 24 is re-used rearing water from the hatchery and is made up primarily of spring water from Indian Ford A Springs originating across the river from the hatchery. This is the same spring water, which is used for the incubation and early rearing of 2m on-station juveniles. In the spring, river water is introduced for acclimation for this pond. The remaining population is reared in Pond 26 which is supplied with spring water from Wonder Springs which is approximately one-half mile downstream and across the river from the main hatchery. These water sources flow naturally into the Klickitat River and make up a part of its total volume; however, they were not historically available as separate spawning/rearing waters.

Wahkiacus Acclimation Facility

This facility will use up to 24 cfs of river water from the Klickitat River. Water quality in the area is acceptable for spring and early summer acclimation. High turbidity during storm events may cause short-term problems in juvenile fish rearing and feeding.

4.2 Indicate risk aversion measures that will be applied to minimize the likelihood for the take of listed natural fish as a result of hatchery water withdrawal, screening, or effluent discharge.

<p>Hatchery water withdrawal</p>	<p><i>Klickitat Hatchery:</i> Water rights total 6000 – 8000 gpm from the gravity intake with another 3,000 pumped from the river. Water rights will be formalized through trust water rights from the Department of Ecology (DOE). Monitoring and measurement of water usage is reported in monthly NPDES reports.</p> <p>Water permits for the Klickitat Hatchery are listed as: S4-01258CWRIS, S428163CWRIS, S4-27553CWRIS,S4-27554CWRIS, S4*07272CWRIS,S3-+22202CWRIS,S4-*07273PWRIS, S4-*07274PWRIS, S4-30084.</p> <p>Lyle Falls Fishway Fishway S4-35252.</p> <p><i>Wahkiacus Acclimation Facility:</i> A new Klickitat River water right permit has been obtained for this site from DOE. S4-34554P</p>
<p>Intake/Screening</p>	<p>Intake structures will be designed and constructed to</p>

<p>Compliance</p>	<p>NMFS specifications at Wahkiacus Acclimation Facility. The Mitchell Act Intake and Screening Assessment (2002) identified the design and alternatives needed to get existing structures at Klickitat Hatchery in compliance with NMFS fish screening standards. From the assessment, YN has been requesting funding for future scoping, design, and construction work of a new intake system.</p>
<p>Hatchery effluent discharges. (Clean Water Act)</p>	<p>Both facilities will operate under the “<i>Federal Aquaculture Facilities and Aquaculture Facilities Located in Indian Country within the Boundaries of the State of Washington</i>” National Pollution Discharge Elimination System (NPDES) general permit (WAG-130021) which conducts effluent monitoring and reporting and operates within the limitations established in its permit administered by the EPA. Monthly and annual reports on water quality sampling, use of chemicals, and compliance records will be available from the EPA.</p> <p>Discharges from the cleaning treatment system are monitored as follows:</p> <p><i>Total Suspended Solids (TSS):</i> Collected 1 to 2 times per month on composite effluent, maximum effluent, and influent samples.</p> <p><i>Settleable Solids (SS):</i> Collected 1 to 2 times per week on effluent and influent samples.</p> <p><i>In-hatchery Water Temperature</i> - Daily maximum and minimum readings.</p> <p>Acclimation ponds will be designed to handle and process effluent during cleaning operations.</p>

Section 5. Facilities

5.1 Broodstock collection facilities (or methods).

Broodstock to be collected at Lyle Falls, Klickitat River Hatchery, and LWSNFH. Adults will be collected throughout the entire run. DNA sampling will be utilized on adults collected in the Klickitat River to ensure that spring Chinook and out-of-basin origin fish are not used as broodstock.

In-basin broodstock adults will be transported by truck to holding facilities at the Klickitat Hatchery. If there is need to augment LWSNFH brood needs, additional adults trapped at Lyle Falls may be transported to Little White Salmon Ponds (Rkm 2.4) of the Big White Salmon River to supply brood need for

USFWS pre-smolt transfers to WAF. Adult transportation protocols defined in Hager and Costello (1999) will be followed.

Time in transport will be less than 1 hour from Lyle to the Klickitat Hatchery and from Lyle to Big White Salmon Ponds.

5.2 Fish transportation equipment (description of pen, tank, truck, or container used).

Adults and juveniles are trucked to holding and acclimation sites.

5.3 Broodstock holding and spawning facilities.

Adult holding ponds at Klickitat will be fed with a combination of river, well and spring water.

See Little White Salmon NFH HGMP for data on adult holding at this facility.

5.4 Incubation facilities.

Incubation facilities at Klickitat hatcheries use well or spring water for incubation. Klickitat Hatchery has 72 stacks of FAL Heath incubators for incubation and hatching. Stack incubators are loaded at 6000-8000 eggs/per tray for hatching. Removal of dead eggs, accurate enumeration, and loadings are adjusted during this time.

See Little White Salmon NFH HGMP for data on incubation facilities at this hatchery.

5.5 Rearing facilities.

Fall Chinook will be reared to volitional release at both the Klickitat Hatchery and Wahkiacus Acclimation Facility.

Ponds (No.)	Pond Type	Volume (cu.ft)	Length (ft.)	Width (ft.)	Depth (ft.)	Dia (ft.)	Flow (gpm)
Klickitat Hatchery							
14	concrete	3500	100	10	3.5		3500

P26	Hypalon Pond	29925	190	45	3.5		5500
Wahkiacus Acclimation							
7	Circulars @ WAF 30' dia. ~6 ft deep	3888			5.5	30	2500

On station Fall Chinook (Klickitat Hatchery 2 million production) fry are ponded in raceways during late February early March. In May the fingerlings are transferred to release Pond 26 at the Klickitat Hatchery until release at approximately 80 fpp.

At Wahkiacus, 2 million pre-smolts will be transferred from LWSNFH in mid-May at ~400 fpp and acclimated until release at 80 fpp in late June.

5.6 Acclimation/release facilities.

The on-station fall Chinook will be released from Pond 26 and reared to release on spring water. The Little White Salmon NFH pre-smolts will be transferred to WAF in mid- May. Once at WAF, these fish will be reared using river water. It should be noted that WAF well water has been developed and may provide beneficial/emergency use at times. Fish will be allowed to migrate volitionally from rearing ponds at both the Klickitat Hatchery and Wahkiacus Acclimation Facility.

As the program is shifted to local broodstock and pre-smolt transfers, eyed-egg and/or green egg transfers from Little White Salmon NFH will be eliminated.

5.7 Describe operational difficulties or disasters that led to significant fish mortality.

Currently, the 4.0 million target release strains facility operations, making best rearing practices difficult to implement. The new Wahkiacus Acclimation Facility will alleviate this problem by increasing rearing capacity. However, there has been no instance of large-scale fish loss for the current program.

5.8 Indicate available back-up systems, and risk aversion measures that will be applied, that minimize the likelihood for the take of listed natural fish that may result from equipment failure, water loss, flooding, disease transmission, or other events that could lead to injury or mortality.

Potential Hazard	Risk Aversion Measure
Equipment failure/Water loss	Both facilities will have multiple water sources available. At Klickitat Hatchery, there is a main river gravity water feed system, three torpedo type river pumps, and several springs available. Backup generator system is automatic in case of power loss. Wahkiacus Acclimation Facility will have both river intake and backup well water source.
Flooding/Water Loss	<p>Both facilities are sited to minimize the risk of catastrophic fish loss from flooding and equipped with low water alarm probes in strategic locations to prevent fish loss due to loss of water. Alarm systems are monitored 24/7 with staff available on-station to respond to problems.</p> <p>At Wahkiacus, under 100-year flood conditions, circular tanks may be overtopped. Fish could be washed out of the system during these events. Debris entering the tanks will need to be removed and tanks and site cleaned and repaired.</p>
Disease Transmission	USFWS fish health guidelines are followed. USFWS fish health pathologists conduct inspections monthly and problems are managed promptly to limit mortality and reduce possible disease transmission.

Section 6. Broodstock Origin and Identity

6.1 Source.

Source is Little White Salmon NFH and the Klickitat River. Broodstock used in the program will be collected at random throughout the run-at-large at Lyle Falls, Klickitat Hatchery, and Little White Salmon NFH.

As the project progresses, adult fall Chinook returning to the Klickitat River will be used for the local on-station release. Klickitat River broodstock will be collected (as needed) for broodstock for the LWSNFH pre-smolt transferred to WAF.

6.2.1 History.

Introductions of fall-run Chinook salmon into the Klickitat River began in 1946 (Marshall et al. 1995), and although a hatchery broodstock was established, tule stocks from various facilities continued until 1986. Beginning in 1986, Klickitat Hatchery production switched from the tule stock to an upriver bright (URB) fall Chinook. Since 2007, 4 million hatchery URB smolts are released on-station annually, primarily for harvest augmentation. Eyed eggs currently are transferred from Little White NFH to the Klickitat Hatchery for final rearing. There is no capture of fall-run Chinook salmon adults at Klickitat Hatchery and eggs are imported yearly from Little White Salmon, or Bonneville hatcheries. Genetic analysis of naturally spawning Klickitat fall Chinook sampled from 1991 to 1994 showed them to be very similar genetically to URB Chinook at Priest Rapids Hatchery and in Hanford Reach and they were closely associated with URB populations at Bonneville and Little White Salmon hatcheries and in the Yakima River (Marshall 2000).

See Little White Salmon Nation Fish Hatchery URB Chinook HGMP for additional egg transfer information.

6.2.2 Annual size.

A total of 4.5 million eyed eggs have been historically transferred to Klickitat Hatchery for this program. The 4.5 million egg-take will continue; however, the program now calls for collecting between 1,250 – 2,500 adults returning to the Klickitat. Adult and jack fall Chinook returning to the Klickitat River will be used as the broodstock source for the program. The broodstock will be collected at the Lyle Falls and Klickitat Hatchery facilities. Approximately 1,250 adults will be needed to produce an on-station release of 2.0 million subyearling fall Chinook. Klickitat River collection (as needed) to support LWSNFH egg take. Broodstock will be collected throughout the entire adult migration period to increase the diversity of life histories being reared at the hatchery. This action will reduce disease risks associated with the importation of eggs from outside of the Subbasin.

6.2.3 Past and proposed level of natural fish in the broodstock.

Estimates of the proportion of wild fish used as broodstock are not known. However, because broodstock were collected at historically from Priest Rapids Hatchery and now Little White Salmon NFH, Klickitat River origin fish have never been used as broodstock.

The new program will use adult fall Chinook that return to the Subbasin as broodstock for the 2.0 million on-station program. Both marked and unmarked fish may be incorporated into the hatchery program. However, because a native run of fall Chinook does not exist above Lyle Falls, the number of natural-origin fish available to the program is limited. Therefore, the program does not establish a goal for the number or proportion of natural-origin fish used as broodstock.

Adult and jack fall Chinook returning to the Klickitat River will be used as the broodstock source for the program. The broodstock will be collected at the Lyle Falls and Klickitat Hatchery. Approximately 1,250 adults will be needed to produce an on-station release of 2.0 million subyearling fall Chinook. Klickitat River collection as needed to support LWSNFH egg take. Broodstock will be collected throughout the entire adult migration period to increase the diversity of life histories being reared at the hatchery.

6.2.4 Genetic or ecological differences.

Tule fall Chinook are not indigenous to the Klickitat Subbasin. Hatchery plants (tule) from outside the Subbasin first occurred in 1946. Releases from the Klickitat Hatchery began in 1952 and continued until 1986. Releases have included stocks from Cowlitz, Toutle, Kalama, Washougal, Bonneville, Cascade, and Ringold hatcheries. The Klickitat fall Chinook program was originally developed to rear tule fall Chinook from the Spring Creek Hatchery. When the Spring Creek program failed to provide the necessary eggs, the program was changed to URB Chinook. The URB Chinook program was intended to provide a better quality fish for the tribal terminal fishery in the lower Klickitat River.

6.2.5 Reasons for choosing.

Using Klickitat River origin fish is expected to increase survival rates, reduce disease transfer risks, and reduce genetic risks to the natural population (HSRG 2005).

6.3 Indicate risk aversion measures that will be applied to minimize the likelihood for adverse genetic or ecological effects to listed natural fish that may occur as a result of broodstock selection practices.

Using native Klickitat River adults for broodstock is expected to minimize straying into adjacent subbasins, thereby reducing genetic risks to ESA -listed Chinook populations in the Columbia and Snake rivers.

Section 7. Broodstock Collection

7.1 Life-history stage to be collected (adults, eggs, or juveniles).

Adults

7.2 Collection or sampling design

Adults will be collected at random throughout the entire upstream migration period. Fish will be collected at Lyle Falls and Klickitat Hatchery.

7.3 Identity.

Klickitat River and URB at Little White Salmon NFH. Program goal to eventually collect all broodstock from the Klickitat River.

7.4 Proposed number to be collected:

A maximum of 2,500 adults will be collected for broodstock.

7.4.1 Program goal (assuming 1:1 sex ratio for adults):

A maximum of 2,500 adults will be collected for broodstock.

7.4.2 Broodstock collection levels for the last twelve years, or for most recent years available.

See Little White Salmon Nation Fish Hatchery URB Chinook HGMP for broodstock use and egg transfer information.

7.5 Disposition of hatchery-origin fish collected in surplus of broodstock needs.

Surplus adults will be distributed to tribal members for ceremonial and subsistence purposes.

7.6 Fish transportation and holding methods.

Fall Chinook captured at Lyle Falls will be transported by truck to Klickitat Hatchery following protocols identified in Hager and Costello (1999). Fall Chinook that swim in to the Klickitat Hatchery will be diverted to adult holding ponds. Adult holding ponds are supplied with a combination of river, well, and spring water. Klickitat River adult transfer to support LWSNFH brood needs may be held at the USFWS White Salmon River Ponds prior to spawning by LWSNFH personnel.

Fall Chinook are collected at the LWS NFH adult ladder. It is generally not required to transport adult fall Chinook. The holding period for upriver bright fall Chinook salmon is very short (about one month as a maximum amount of time). The goal for all species is to achieve a 2.5% or less pre-spawning mortality rate during the holding period.

7.7 Describe fish health maintenance and sanitation procedures

applied.

USFWS fish health protocols will be followed for adult holding.

7.8 Disposition of carcasses.

Carcasses of fall Chinook spawned through the programs will be buried at a local landfill. If they receive certification by USFWS, they may be planted as part of a nutrient enhancement project in the Klickitat River Subbasin.

7.9 Indicate risk aversion measures that will be applied to minimize the likelihood for adverse genetic or ecological effects to listed natural fish resulting from the broodstock collection program.

All trapped listed species are identified, quickly sampled and returned to the river. Trapping facilities are designed to NMFS criteria to minimize handling stress and mortality.

Converting to local broodstock is expected to reduce adult straying rates to other subbasins, thereby decreasing genetic risks to other Chinook populations.

Section 8. Mating

8.1 Selection method.

The spawning protocol mandates the use of a spawning population of at least 500 adults. Spawners are selected and mated randomly from the population maintained in the hatchery holding pond. Fish are spawned throughout the entire run to help ensure that the run timing for the stock is maintained.

8.2 Males.

Jacks will be used in proportion to that observed in the natural run.

8.3 Fertilization.

A 1:1 male to female ratio will be used for fertilization.

8.4 Cryo-preserved gametes.

Cryo-preserved gametes are not used.

8.5 Indicate risk aversion measures that will be applied to minimize the likelihood for adverse genetic or ecological effects to listed natural fish resulting from the mating scheme.

DNA samples will be collected on all adult fish collected for broodstock. We will attempt to use only fish which type similar to mid-Columbia Bright (MCB) fall Chinook populations for brood stock. Over time we expect the local (Klickitat) brood source may develop a unique DNA profile and these fish would be incorporated into brood stock to the maximum extent possible.

No listed fish will be spawned or mated as part of this program.

Section 9. Incubation and Rearing.

9.1.1 Number of eggs taken and survival rates to eye-up and/or ponding.

Eggs are taken from fish spawned at Little White NFH. Once “eyed”, 4.5 million eggs will be transferred to Klickitat Hatchery. Information in this table reflect the total numbers of URB eggs spawned at that respective facility, not at the Klickitat Hatchery. Percentages reflect survival at the Klickitat Hatchery.

Year	Egg Take	Eyed-Ponding Survival (%)
1995	17,345,900	98.10
1996	14,533,500	99.13
1997	17,007,000	99.24
1998	13,981,300	98.09
1999	16,089,600	98.72
2000	15,349,500	97.89
2001	13,389,500	94.28
2002	13,732,550	99.54
2003	13,820,500	98.90
2004	12,753,500	92.30
2005	13,500,100	92.33
2006	14,412,102	91.38
2007	11,338,617	91.46
2008	9,158,431	89.13
2009	7,462,632	85.49
2010	8,901,550	87.84
2011	9,158,431	92.40
2012	8,250,700	87.92

At Klickitat Hatchery, eyed egg to ponding survival is approximately 98.2%. Some green eyed shipping loss was experienced with 2 early takes of URB eggs from LWS, BY 2010.

9.1.2 Cause for, and disposition of surplus egg takes.

Variability in fecundity and egg survival may result in surplus eggs being collected. Surplus eggs may be released or destroyed dependent on the results

of consultations with the co-managers.

9.1.3 Loading densities applied during incubation.

Klickitat Hatchery has 72 stacks of FAL Heath incubators for incubation and hatching. Stack incubators are loaded at 6000-8000 eggs/per tray for hatching. Removal of dead eggs, accurate enumeration and loadings are adjusted during this time. See section 5.4 for load and hatching criteria. Integrated Hatchery Operations Team (IHOT) species-specific incubation recommendations are followed for water quality, flows, temperature, substrate and incubator capacities.

9.1.4 Incubation conditions.

Integrated Hatchery Operations Team (IHOT) species-specific incubation recommendations will be followed for water quality, flows, temperature, substrate, and incubator capacities at Klickitat hatchery.

Harmful silt and sediment is cleaned from incubation systems regularly while eggs are monitored to determine fertilization and mortality. Incubation water is from Indian Ford A Springs located across the river from the hatchery. Temperature is monitored by thermograph and recorded; temperature units (TU) are tracked for embryonic development. Dissolved oxygen content is monitored and has been at acceptable levels of saturation with a minimum criteria of 8 parts per million (ppm). When using artificial substrate, vexar, or bio-rings, egg densities within incubation units are reduced by 10%.

9.1.5 Ponding.

Fall Chinook fry are transferred from Heath trays for ponding upon button-up and swim-up. Fry are ponded when: a visual inspection of the amount of yolk sac remaining with the yolk slit closed to approximately 1 millimeter wide (approximately 1,600 – 1,800 temperature units) or based on (95% yolk absorption) KD factor. The mean weight for fry ponded is 700-800 fpp. At this time, fry are transferred to raceways for rearing.

9.1.6 Fish health maintenance and monitoring.

USFWS fish health guidelines are followed. Hatchery staff conducts daily inspection, visual monitoring and sampling from eye, fry fingerling and sub-yearling stages. Potential problems are immediately communicated to the USFWS fish health specialist. In addition, fish health specialists conduct inspections monthly. Potential problems are managed promptly to limit mortality and reduce possible disease transmission. At spawning, eggs are water-hardened in iodophor as a disinfectant. Formalin (37% formaldehyde) is periodically dispensed into water supplied to the incubators and raceways to control fungus growth on eggs. Formalin may also be used on parasite loads on juvenile salmon, if recommended by a fish health specialist. Treatment dosage and duration varies by life-stage and condition being treated. All fish disease control procedures are conducted consistent with USFWS Policy for fish reared in the Klickitat and Little White Salmon National Fish Hatchery.

9.1.7 Indicate risk aversion measures that will be applied to minimize the likelihood for adverse genetic and ecological effects to listed fish during incubation.

Listed fish are not incubated for this program.

9.2.1 Provide survival rate data (*average program performance*) by hatchery life stage (fry to fingerling; fingerling to smolt) for the most recent twelve years, or for years dependable data are available.

Year	Fry-fingerling Survival (%)
1995	96.50
1996	98.86
1997	98.78
1998	99.19
1999	99.40
2000	96.13
2001	98.69
2002	97.72
2003	98.43
2004	94.21
2005	99.15
2006	89.16
2007	98.04
2008	98.32
2009	97.08
2010	99.89
2011	99.01
2012	99.00

At Klickitat Hatchery, eyed egg to smolt release survival for the last nine years has averaged 91%.

9.2.2 Density and loading criteria (goals and actual levels).

The pond loading densities maintained at the Klickitat Hatchery are consistent with those recommended by Piper et al. (1982; 6 lb/gpm and 0.75 lb/ft³) and Banks (1994; 0.125 lb/ft³/in) (BAMP 1998).

Fry are transferred from the Heath incubation trays to vinyl raceways for start feeding and continued rearing. The raceways have flow through water circulation.

9.2.3 Fish rearing conditions.

Fish are to be reared on a combination of river, spring and well water at Klickitat Hatchery and WAF. Information provided below is for the Klickitat Hatchery, but will apply rearing at the WAF as well once this facility is constructed.

Temperature, dissolved oxygen and pond turn over rate are monitored. IHOT standards are followed for: water quality, alarm systems, predator control measures (netting) to provide the necessary security for the cultured stock, fish loadings and densities. Settleable solids, unused feed, and feces are removed regularly to ensure proper cleanliness of rearing containers. All ponds are broom-cleaned as needed and pressure-washed between broods. Temperature and dissolved oxygen are monitored and recorded daily during fish-rearing. Temperatures during the rearing cycle range from a high of 65 to a low of 33 degrees F. Raceway vessels are cleaned on an as-needed basis. Netting covers are placed over acclimation rearing ponds to minimize predation.

9.2.4 Indicate biweekly or monthly fish growth information (average program performance), including length, weight, and condition factor data collected during rearing, if available.

Table 9.2.4.1. Growth data for Little White Salmon National Fish Hatchery fall Chinook during final rearing at the Klickitat Hatchery, 2010.			
Rearing Period	Length (mm)	Weight (fpp)	Condition Factor
02/10	Na	Na	Na
03/10	Na	890	Na
04/10	Na	359	Na
05/10	Na	159	Na
06/10	82	73	1.114

9.2.5 Indicate monthly fish growth rate and energy reserve data (average program performance), if available.

Table 9.2.5.1. Growth data for Little White Salmon National Fish Hatchery fall Chinook during final rearing at the Klickitat Hatchery, 2010.			
Rearing Period	Length (mm)	Weight (fpp)	Condition Factor
02/10	Na	Na	Na
03/10	Na	890	Na
04/10	Na	359	Na
05/10	Na	159	Na

06/10	82	73	1.114
-------	----	----	-------

9.2.6 Indicate food type used, daily application schedule, feeding rate range (e.g. % B.W./day and lbs/gpm inflow), and estimates of total food conversion efficiency during rearing (average program performance).

Table 9.2.6.1. Feed data for Little White Salmon National Fish Hatchery fall Chinook during rearing at the Klickitat Hatchery, 2010.

Rearing Period	Food Type	Application Schedule (# feedings/day)	Feeding Rate Range (%B.W./day)	Lbs. Fed per gpm of Inflow	Food Conversion During Period
3/15/10 – 4/30/2010	Ewos Micro#1 & 2	4-8	1.50-2.00	0.023	0.48
4/30/2010 – 6/14/2010	Ewos Pacific 1.2mm	2-4	1.5	0.026	0.55
6/14/2010 – 6/28/2010	Ewos Vita 1.5 mm	2	1.5	0.051	0.54

9.2.7 Fish health monitoring, disease treatment, and sanitation procedures.

Fish Health Monitoring	A fish health specialist inspects fish monthly and checks both healthy and if present symptomatic fish. Based visual detection of pathological problems, age of fish, and the history of the facility, the pathologist determines the appropriate tests. External signs such as lesions, discolorations, and fungal growths will lead to internal examinations of skin, gills, and organs. Kidney and spleen are checked for bacterial kidney disease (BKD). Blood is checked for signs of anemia or other pathogens. Additional tests for virus or parasites are done if warranted.
Disease Treatment	Appropriate therapeutic treatment will be prescribed to control and prevent further outbreaks. Dead fish are collected and disposed of at a landfill. Fish health and or treatment reports are kept on file.
Sanitation	All eggs brought to the facility are surface-disinfected with iodophor (per disease policy). All equipment (nets, tanks, boots, etc.) is disinfected with iodophor between different fish/egg lots. Different fish/egg lots are physically isolated from each other by separate ponds or incubation units. The intent of these activities is to prevent the horizontal spread of pathogens by splashing water. Tank trucks are disinfected between the hauling of adult and juvenile fish. Foot baths containing disinfectant are strategically located on the hatchery grounds to prevent spread of pathogens.

9.2.8 Smolt development indices (e.g. gill ATPase activity), if applicable.

Fall Chinook are released in May-June as sub-yearling smolts. Program goal has been to release fish when they reach 80 fpp. Along with size, appearance, and release time are used to indicate the readiness of the population for emigration.

ATPase data will be collected on fish released at Klickitat Hatchery and the Wahkiacus Acclimation Facility to determine smoltification status.

9.2.9 Indicate the use of "natural" rearing methods as applied in the program.

None.

9.2.10 Indicate risk aversion measures that will be applied to minimize the likelihood for adverse genetic and ecological effects to listed fish under propagation.

Listed fish are not under propagation.

Section 10. Release

10.1 Proposed fish release levels.

On-station at the Klickitat Hatchery (2.0 million) smolts at 50-80 fish per pound (FPP).

LWSNFH pre-smolt transfers to WAF (2.0 million) smolts at 50-80 fish per pound (FPP).

10.2 Specific location(s) of proposed release(s).

Fall Chinook will be released volitionally from rearing sites located at RKm 27 (Wahkiacus Hatchery). The remaining Fall Chinook production (2,000,000) will be released volitionally from the Klickitat Hatchery (RKm 68).

10.3 Actual numbers and sizes of fish released by age class through the program.

Release Year	Fingerling Release		
	No.	Date (MM/DD)	Avg Size (fpp)
1996	4,380,000	05/16-06/08	64.0
1997	3,625,870	May	65.0
1998	4,387,480	05/21-05/31	71.0
1999	4,289,100	06/02-06/07	71.0
2000	3,972,500	05/15-05/22	55.0
2001	3,850,300	05/22-05/25	66.0
2002	3,968,900	06/03-06/07	65.0
2003	3,664,100/	6/03-6/19/	73.0
	520,000	7/16-7/20	79.0
2004	2,590,650/	6/14-18/	62.2
	1,635,000	7/6-13/	69.0
2005	2,150,500/	6/12-16/	77.0
	2,397,800	6/19-21/	69.6
2006	4,548,270	6/12-6/16	71.8
2007	2,158,500	6/25-29/	83.1
	2,198,000	6/11-15/	82.8
2008	1,539,477	6/16-20/	75
	2,012,960	6/26-30/	69
2009	2,018,200	6/12-14/	85
	2,357,900	6/9-11/	74
2010	2,073,311	7/7-10/	75
	2,079,425	6/28-7/2	67.9
2011	1,987,240	6/6-9/	88
	1,988,937	6/1-7/	92
2012	1,584,020	6/18-21/	70
	1,856,575	6/18-21/	70

10.4 Actual dates of release and description of release protocols.

Fish will be allowed to migrate volitionally from the rearing ponds. Fish that do not migrate volitionally from the ponds will be seined and destroyed. Carcasses will be buried in an upland landfill.

10.5 Fish transportation procedures, if applicable.

Fingerlings are not transported.

10.6 Acclimation procedures (*methods applied and length of time*).

Chinook for this program will be reared on a combination of Klickitat River, adjacent spring water or well water for prior to release. Rearing on parent river water, or acclimation for several weeks to parent river water, is done to ensure strong homing to the release site, thus reducing the stray rate to natural populations.

10.7 Marks applied, and proportions of the total hatchery population marked, to identify hatchery adults.

1) 16.5% of all fish released will be adipose-clipped and coded-wire-tagged (CWT).

2) with infrastructure improvements to allow for safe marking, it is intended that 100% of all will be adipose-clipped until these fish/egg transfers are eliminated.

3) Klickitat River origin juveniles will be tagged with a blank cwt and ad-clipped.

10.8 Disposition plans for fish identified at the time of release as surplus to programmed or approved levels

N/A.

10.9 Fish health certification procedures applied pre-release.

Prior to release from Wahkiacus and Klickitat Hatcheries the population health and condition is established by the USFWS Fish Health Pathologist. This is commonly done 1-3 weeks pre-transfer and up to 6 weeks on systems with pathogen-free water and little or no history of disease. Prior to this examination, whenever abnormal behavior or mortality is observed, staff contacts the USFWS Fish Health Pathologist. The pathologist examines affected fish and recommends appropriate treatment. Reporting and control of selected fish pathogens are done in accordance with the Co-managers' Fish Disease Control Policy and USFWS guidelines.

10.10 Emergency release procedures in response to flooding or water system failure.

Emergency procedures and disposition of fish will adhere to the protocols and procedures set forth in approved operation plans. If the program is threatened by ecological or mechanical events, the Complex Manager will contact YN fish management. If an on-station emergency release is authorized, personnel will pull screens and sumps and fish will be force- released into the Klickitat River.

10.11 Indicate risk aversion measures that will be applied to minimize the likelihood for adverse genetic and ecological effects to listed fish resulting from fish releases.

- Fish are reared to sufficient size (80 fpp) so that smoltification occurs simultaneously within almost the entire population. This will reduce retention in the streams after release.
- Rearing on natal river water or acclimation for several weeks to this river

water is done to ensure strong homing to the hatchery, thus reducing adult stray rate to streams outside of the Klickitat River basin.

- The Yakama Nation transition plan calls for moving ½ the program (2 million) to a lower Klickitat River acclimation site, assuming funding for construction and O&M of the Wahkiacus Acclimation Facility is secured. . This will reduce interactions with listed steelhead in the Klickitat River Subbasin.

Section 11. Monitoring and Evaluation of Performance Indicators

11.1.1 Describe plans and methods proposed to collect data necessary to respond to each "Performance Indicator" identified for the program.

Harvest: Fisheries will be monitored by the co-managers to determine harvest rates and numbers in Columbia River and Klickitat River Subbasin fisheries. Ocean fisheries will be monitored by the WDFW and other entities by sampling fishing boats and fishers when they return to ports or to fish processing stations.

Smolt-to-adult Survival Rate: A portion of all releases will be marked with CWT and adipose fin-clipped. These tags will be recovered from fish caught in all fisheries, spawning ground surveys, hatchery returns, and from tags voluntarily submitted by the public.

Adult Straying: Regional M&E efforts will be used to track the number and capture location of fish originating from the Klickitat River. Klickitat River fish will be tagged with a wire-tag so that they can be identified in fisheries, in spawning ground surveys, and at hatcheries.

DNA (Broodstock composition): A portion of the fin collected from adult fall Chinook at broodstock collection facilities will be taken and the DNA analyzed. The adults will be given a unique numeric mark and then released to adult holding facilities. Only those adults identified as originating from the Klickitat River Subbasin will be used as broodstock.

Juvenile Health Monitoring: Juvenile fish at Klickitat Hatchery are monitored on a routine basis by the hatchery staff to determine the condition of fry, fingerlings, and smolts. Samples will be taken by the USWS Fish Health pathologist to determine the health of fry, fingerling, and smolts prior to release. Sampling of fingerlings for tag retention and fin mark quality, prior to release, is conducted by YN marking program.

Environmental Monitoring: Environmental monitoring is conducted at hatchery facilities to ensure that the facilities meet the requirements of the National Pollution Discharge Elimination System (NPDES) permit and is also used in managing fish health. On a short-term basis, environmental monitoring helps

identify when changes to hatchery practices are required. The following parameters are currently monitored -

- Total Suspended Solids (TSS)
- Settleable Solids (SS)
- Water Temperature

Water Diversion Screens: Facility water intakes will be designed in accordance with NMFS standards. Velocity measurements will be taken at each screening system each year to ensure the criteria are being met. Screens will also be inspected each year for any problems (missing panels, debris, etc.) and corrective actions taken.

Hatchery Operations: An annual report will be written documenting program operations, disease problems, treatment, broodstock collection, number of fish released, fish size, and release date.

11.1.2 Indicate whether funding, staffing, and other support logistics are available or committed to allow implementation of the monitoring and evaluation program.

BPA and Mitchell Act funding are expected to be sufficient to implement program and associated monitoring.

11.2 Indicate risk aversion measures that will be applied to minimize the likelihood for adverse genetic and ecological effects to listed fish resulting from monitoring and evaluation activities.

- For fall Chinook, spawning/carcass surveys occur at a time when adult steelhead are not spawning.
- Any disturbance of adult or juvenile steelhead during spawning and carcass surveys is expected to be minimal because surveys are performed only once per week within a given river reach.
- All trapped listed species are identified, quickly sampled and returned to the river. Trapping facilities are designed to NMFS criteria to minimize handling stress and mortality.

Section 12. Research

12.1 Objective or purpose.

No research activities are proposed for fall Chinook that will affect listed fish populations in the Klickitat River, other than monitoring and evaluation activities described above and in Appendix 1.

Section 13. Attachments and Citations

13.1 Attachments and Citations

Banks, J.L. 1994. Raceway Density and Water flow as Factors Affecting Spring Chinook Salmon (*Oncorhynchus tshawytscha*) during Rearing and after Release. *Aquaculture*, 119 (1994) 201-217. 54: pp. 137-147.

Biological Assessment and Management Plan (BAMP). 1998. Mid-Columbia River Hatchery Program. National Marine Fisheries Service, U.S. Fish and Wildlife Service, Washington Department of Fish and Wildlife, Confederated Tribes of the Yakama Indian Nation, Confederated Tribes of the Colville Indian Reservation, and the Confederated Tribes of the Umatilla Indian Reservation. Mid-Columbia Mainstem Conservation Plan.

Busack, Craig, Kenneth Currens, Todd Pearsons, Lars Moberg. 2005. "Tools for Evaluating Ecological and Genetic Risks in Hatchery Programs", 2004 Final Report, Project No. 200305800, 91 electronic pages, (BPA Report DOE/BP-00016399-1).

Byrne, J., R. McPeak, B. McNamara - Washington Department of Fish and Wildlife. 2001. Bull Trout Assessments in the Columbia River Gorge, FY-2000 Annual Report, Report to Bonneville Power Administration, Contract No. 00000651, Project No. 199802600, 85 electronic pages (BPA Report DOE/BP-00000651-1).

Dawley, E. M., R.D. Ledgerwood, T.H Blahm, R.A. Kirn, and A.E. Rankis. 1984. Migrational Characteristics And Survival Of Juvenile Salmonids entering the Columbia River estuary During 1983. Annual Report to the Bonneville Power Administration, Portland, OR.

Flagg, T.A., B.A. Berejikian, J.E. Colt, W.W. Dickhoff, L.W. Harrell, D.J. Maynard, C.E. Nash, M.S. Strom, R.N. Iwamoto, and C.V.W. Mahnken. 2000. Ecological and behavioral impacts of artificial production strategies on the abundance of wild salmon populations. U.S. Dept. Commer., NOAA Tech. Memo. NMFS-NWFSC-41: 92pp.

Ford, M.J., T. Cooney, P. McElhany, N. Sands, L. Weitkamp, J. Hard, M. McClure, R. Kope, J. Myers, A. Albaugh, K. Barnas, D. Teel, P. Moran, and J. Cowen. 2010. Status review update for Pacific salmon and steelhead listed under the Endangered Species Act: Northwest. Draft U.S. Department of Commerce, NOAA Technical Memorandum.

Gray, S.W. 2007. Determine the Origin, Movements, and Relative Abundance of Bull Trout in Bonneville Reservoir. Available at:
<https://pisces.bpa.gov/release/documents/documentviewer.aspx?doc=P103326>

Hager, R.C. and R.J. Costello. 1999. Optimal Conventional and Semi-natural Treatments for the Upper Yakima Spring Chinook Salmon Supplementation

Project: Treatment Definitions and Descriptions and Biological Specifications for Facility Design. Proj. No. 95-06404, Prepared for Bonneville Power Administration. Portland, OR.

Hawkins, S.W., Tipping, J. M. 1999. Predation By Juvenile Hatchery Salmonids on Wild Fall Chinook Salmon Fry in the Lewis River, Washington. *California Fish and Game* **85**(3):124-129

Hatchery Scientific Review Group (HSRG). 2005. Hatchery Reform in Washington State: Principles and Emerging Issues. *Fisheries*. Volume 30, Number 6. June 2005.

Hatchery Scientific Review Group (HSRG). 2009. System-Wide Report on Columbia River Basin Hatchery Reform. Available at: http://hatcheryreform.us/hrp/reports/system/welcome_show.action

Harza. 1998 The 1997 and 1998 technical study reports, Cowlitz River Hydroelectric Project. Vol. 2: 35-42.

Howell, P., K. Jones, D. Scarnecchia, L. LaVoy, W. Knedra and D. Orrman. 1985. Stock assessment of Columbia River anadromous salmonids. Vol: 1. U.S. Dep. Energy, Bonneville Power Administration. Project No. 83-335, 558 pp.

IHOT (Integrated Hatchery Operations Team). 1995. Operation plans for anadromous fish production facilities in the Columbia River basin. Volume III-Washington. Annual Report 1995. Bonneville Power Administration, Portland Or. Project Number 92-043. 536 pp.

Interior Columbia Basin Technical Review Team (ICTRT). 2007. Viability Criteria for Application to Interior Columbia Basin Salmonid ESUs (Review Draft March 2007). Prepared by the Interior Columbia Basin Technical Recovery Team. Portland, OR, and Seattle, WA.

Recovery Plan for the Klickitat Population of the Middle Columbia River Steelhead Distinct Population Segment (NOAA-Portland 2009).

Marshall, A.R. 2000. Genetic analysis of Chinook populations in the Klickitat River. Unpublished draft report to Yakima Klickitat Fisheries Project, WDFW Genetics Unit. Olympia, WA.

Marshall, A.R., Smith, C., Brix, R., Dammers, W., Hymer, J., and LaVoy, L. 1995. Genetic diversity units and major ancestral lineages for Chinook salmon in Washington. Edited by C. Busack and J.B. Shaklee. *Wash. Dep. Fish Wildl. Tech. Rep. RAD 95 02*. Olympia, Wash. pp. 111–173.

Narum S. R., M. Powell, R. Evenson, B. Sharpe and A. Talbot. 2006. Microsatellites Reveal Population Substructure of Klickitat River Native Steelhead and Genetic Divergence from an Introduced Stock. *North American Journal of Fisheries Management* **26**:147-155.

NMFS (National Marine Fisheries Service). 1999. Biological opinion on artificial propagation in the Columbia River Basin.

NMFS (National Marine Fisheries Service). 2008. Anadromous Fish Passage Facility Design. Available at: http://www.nwr.noaa.gov/Salmon-Hydropower/FERC/upload/Fish_Passage_Design.pdf

NMFS (National Marine Fisheries Service). 1999. Biological opinion on artificial propagation in the Columbia River Basin.

NMFS. 2009. Middle Columbia River Steelhead Distinct Population Segment ESA Recovery Plan. Corrected Version November 30, 2009. Available at <http://www.nwr.noaa.gov/Salmon-Recovery-Planning/Recovery-Domains/Interior-Columbia/Mid-Columbia/Mid-Col-Plan.cfm> (accessed December 16, 2010).

NOAA. 2005. Updated Status of Federally Listed ESU's of West Coast Salmon and Steelhead. NOAA Technical Memorandum NMFS-NWFSC-66.

NOAA. 2007. Environmental Assessment of NOAA's National Marine Fisheries Service's (NMFS) approval of Five Fisheries Management and Evaluation Plans For Tributaries of the middle Columbia River Submitted by the Oregon Department of Fish and Wildlife (ODFW) and Washington Department of Fish And Wildlife (WDFW), and of NMFS' Determination that the Plans Adequately Address Section 4(d) Limit 4 Criteria and Do Not Appreciably Reduce the Likelihood of Survival and Recovery of Salmon and Steelhead Listed Under the Endangered Species Act

NPPC. 2004. Klickitat Subbasin Plan. Prepared for the Northwest Power and Conservation Council. Prepared by the Yakama Nation, Klickitat County, and Washington Department of Fish and Wildlife.

Piper, R. et al. 1982. Fish Hatchery Management. United States Dept. of Interior, Fish and Wildlife Service. Washington, D.C.

Seidel, P.. 1983. Spawning Guidelines for Washington Department of Fish and Wildlife Hatcheries. Washington Department of Fish and Wildlife. Olympia, WA.

Thiesfeld, S.L., R.H. McPeak, B.S. McNamara, and I. Honanie. 2002. Bull trout population assessment in the White Salmon and Klickitat Rivers, Columbia River Gorge, Washington. Fiscal Year 2001 Annual Report. BPA Contract # 00004474-00001, Project # 1999-024-00.

USFWS (U.S. Fish and Wildlife Service). 1994. Biological assessment for operation of U.S. Fish and Wildlife Service operated or funded hatcheries in the Columbia River Basin in 1995-1998. Submitted to National Marine Fisheries Service (NMFS) under cover letter, dated August 2, 1994, from William F. Shake, Acting USFWS Regional Director, to Brian Brown, NMFS.

_____. 1998. Biological Opinion for the Effects to Bull Trout from Continued Implementation of Land and Resource Management Plans and Resource Management Plans as Amended by the Interim Strategy for Managing Fish-Producing Watersheds in Eastern Oregon and Washington, Idaho, Western Montana, and Portions of Nevada (INFISH) and the Interim Strategy for Managing Anadromous Fish-Producing Watersheds in Eastern Oregon and Washington, Idaho, and Portions of California (PACFISH). Region 1, Portland, OR.

Wahkiacus Hatchery Conceptual Design Study 2010. Yakima/Klickitat Fisheries Program. Toppenish, WA

Washington Department of Fisheries. 1991. Stock Transfer Guidelines. Hatcheries Program, Washington Department of Fisheries. Olympia, WA.

Washington Department of Fish and Wildlife. Fisheries Management and Evaluation Plan WDFW, 2003. Lower Columbia River. Submitted to NMFS. Portland,OR.

Washington State Conservation Commission. 1999. Salmonid habitat limiting factors water resource inventory area 30 - Klickitat watershed. Final Report.

WDFW and NWIFC 1998. Salmonid Disease Control Policy of the Fisheries Co-Managers of Washington State.

Witty, K., C. Willis, and S. Cramer. 1995. A review of potential impacts of hatchery fish on naturally produced salmonids in the migration corridor of the Snake and Columbia Rivers. S.P. Cramer and Associates, Inc., Gresham, OR.

Yakama Nation 2012. *Draft* Klickitat Anadromous Fisheries Master Plan, Yakima/Klickitat Fisheries Program. Toppenish, WA.

Zendt, J., N. Romero, S. Keep, and M. Babcock. 2013. Klickitat Subbasin Monitoring and Evaluation - Yakima/Klickitat Fisheries Project, 2010-2012 Annual Report. BPA Document ID # P132813.
<https://pisces.bpa.gov/release/documents/documentviewer.aspx?doc=P132813>

Section 14. CERTIFICATION LANGUAGE AND SIGNATURE OF RESPONSIBLE PARTY

14.1 Certification Language and Signature of Responsible Party

“I hereby certify that the information provided is complete, true and correct to the best of my knowledge and belief. I understand that the information provided in this HGMP is submitted for the purpose of receiving limits from take prohibitions specified under the Endangered Species Act of 1973 (16 U.S.C.1531-1543) and regulations promulgated thereafter for the proposed hatchery program, and that any false statement may subject me to the criminal penalties of 18 U.S.C. 1001, or penalties provided under the Endangered Species Act of 1973.”

Name, Title, and Signature of Applicant:

Certified by _____ Date: _____

ADDENDUM A. PROGRAM EFFECTS ON OTHER (AQUATIC OR TERRESTRIAL) ESA-LISTED POPULATIONS

15.1) List all ESA permits or authorizations for USFWS ESA-listed, proposed, and candidate salmonid and non-salmonid species associated with the hatchery program.

No permits in place for this new program. They will be developed through consultation with appropriate agencies as facilities and programs are developed.

15.2) Describe USFWS ESA-listed, proposed, and candidate salmonid and non-salmonid species and habitat that may be affected by hatchery program.

Hatchery operations may impact USFWS listed Klickitat River bull Trout (*Salvelinus confluentus*). Bull trout are listed as Threatened by the USFWS. The USFWS has designated the West Fork Klickitat River and Klickitat River reaches adjacent to the Yakama Indian Reservation as Critical Habitat (Federal Register 2005). Stream habitat in the Klickitat River Basin has been impacted by human activities associated with agriculture, logging, recreation, and urban development.

Hatchery facilities are located both within and near the Klickitat River. Water for rearing anadromous fish at the Klickitat River hatchery is diverted from the river. New juvenile acclimation sites are being developed at Wahkiacus Acclimation Facility that will disturb upland and riparian habitat near the stream channel. A diversion structure or mobile system will also be built at this facility to provide water for acclimating hatchery smolts.

Other listed or candidate species that may be impacted by the construction and operation of the Wahkiacus Acclimation Facility include:

Oregon Spotted Frog (<i>Rana pretiosa</i>)	Candidate
Bald Eagle (<i>Haliaeetus leucocephalus</i>)	Threatened
Northern Spotted Owl (<i>Strix occidentalis</i>)	Threatened

Possible impacts to these species from construction or operation of new facilities have not been quantified.

15.3) Analyze effects.

Bull Trout

Possible hatchery operational effects to listed bull trout in the Klickitat River are described below. The effects are expected to be on-going while the hatchery program remains in place.

Water diversion: Water is diverted from the Klickitat River for hatchery operations. This results in a decrease in the amount and quality of stream habitat at both the Wahkiacus (~0.1 miles) and Klickitat River (~0.25 miles). The loss in habitat could result in a decrease in bull trout abundance. However, because bull trout are primarily found in the West Fork Klickitat River and

tributaries higher in the Subbasin than the hatchery locations, impacts to bull trout are assumed minor.

Diversions Screens:

Klickitat River Hatchery: The Mitchell Act Intake and Screening Assessment (2002) identified design and alternatives needed to get existing diversion and screening structures in compliance with NOAA fish screening standards. From the assessment, YN has requested funding for future scoping, design, and construction work of a new intake system. Staff has not reported any bull trout entrained into hatchery facilities or impinged on screen surfaces.

Wahkiacus Acclimation Facility: This facility will be equipped with screens that meet NMFS fry screening criteria.

Waste and Pollutants: Both facilities will operate under the “Upland Fin-Fish Hatching and Rearing” National Pollution Discharge Elimination System (NPDES) general permit which conducts effluent monitoring and reporting and operates within the limitations established in its permit administered by the Washington Department of Ecology (DOE). The limitations listed in the permit are assumed to be protective of water quality and therefore the hatchery waste water is likely to have little impact on bull trout.

Disease: Outbreaks in the hatcheries may cause significant adult, egg, or juvenile mortality. Over the years, advances in rearing densities, disease prevention and fish health monitoring have greatly improved the health of the programs at Klickitat Hatchery. Policies and Procedures for Columbia Basin Anadromous Salmonid Hatcheries (IHOT 1995) Chapter 5 have been instrumental in reducing disease outbreaks. Fish are planted and transferred after a fish health specialist has determined the populations’ health. The level of indirect take of bull trout from disease is unknown.

Broodstock Collection: Hatchery broodstock will be collected at traps located at Lyle Falls and Klickitat River Hatchery. Although no bull trout have been collected in the past, operation of the new trapping facilities at Lyle falls could result in the capture and handling of both bull trout juveniles and adults. As the facility is designed to NMFS criteria, impacts to bull trout are expected to be minor. Any bull trout captured will be released unharmed to the stream.

Acclimation Facilities: New acclimation facilities are to be constructed at Wahkiacus Acclimation Facility located at Rkm 27. The diversion structure will be screened to meet NMFS criteria for fry. Impacts to bull trout from the diversion structure are expected to be minimal.

Release of Juveniles: The program will release 4.0 million fall Chinook smolts, at a size range of 80-100 mm, into the Klickitat River each year. If it assumed that Chinook can consume fish that are up to 33% of their body length, there is the possibility that bull trout less than 36 mm may be susceptible to predation (See section 2). Because fall Chinook smolts will not be released in the primary bull trout spawning stream (West Fork Klickitat River) it is unlikely that the hatchery smolts will prey on, or compete with, listed bull trout.

Food: The carcasses of returning hatchery fall Chinook adults will increase stream productivity which should result in an increase in food abundance for bull trout. Additionally, fall Chinook juveniles (both hatchery and wild) could provide a food source for adult bull trout.

Monitoring and Evaluation: Smolt trapping may be used to determine if hatchery fall Chinook migrate quickly through the system after release. Some bull trout may be captured and handled at the trapping facilities; these fish will be released unharmed to the stream.

Oregon Spotted Frog

Neither hatchery operations nor proposed new facilities are likely to adversely impact this species. The only known population of Oregon Spotted Frog in the Klickitat River Subbasin is located in the Conboy Lake National Wildlife Refuge (NWR) managed by USFWS (Klickitat Subbasin Plan 2004). The refuge is located approximately 10 miles east of Trout Lake and 7 miles southwest of Glenwood in the Glenwood Valley/Camas Prairie area.

Bald Eagle

Bald eagles can be found throughout the year in the Klickitat River Subbasin. Because this species feeds on salmon, increased hatchery production should result in an increase in food for this species as a result of more adult fish returning to the Subbasin. Bald eagle surveys will be conducted prior to constructing any new facilities in the Subbasin.

Northern Spotted Owl

No impacts are expected to spotted owls because the existing facilities and the proposed facilities are not located in spotted owl habitat.

15.4) Actions taken to minimize potential effects.

Bull trout

Diversion Screens: All intake screens will be updated to meet NMFS screen criteria for fry.

Waste and Pollutants: All terms associated with the NPDES Permit will be implemented and followed.

Broodstock Collection: Any juvenile or adult bull trout captured during broodstock collection activities will be returned safely to the stream channel. Trapping facilities will be designed to meet NMFS standards.

Acclimation Facilities: These facilities will be sited to reduce impacts to riparian and stream habitats to the extent possible. The YN will coordinate the location and construction of this facility with USFWS staff to minimize or avoid impacts to all listed species.

Monitoring and Evaluation: Bull trout collected during juvenile trapping operations will be released unharmed to the stream.

Oregon Spotted Frog

Prior to constructing any facility, stream and riparian areas near proposed sites will be surveyed for the presence Oregon Spotted Frogs. If this species is found, the YN will

coordinate with USFWS staff to develop mitigation and protection measures. This activity will be further in the EIS required for a Step 2 project.

Bald Eagle

Acclimation facilities will not be located near bald eagle nests.

Northern Spotted Owl

No activities are proposed that will impact this species because no facilities or activities are planned for areas inhabited by spotted owls or which are suitable spotted owl habitat.

15.5) References

IHOT (Integrated Hatchery Operations Team). 1995. Operation plans for anadromous fish production facilities in the Columbia River basin. Volume III-Washington. Annual Report 1995. Bonneville Power Administration, Portland Or. Project Number 92-043. 536 pp.

NPPC 2004. Klickitat Subbasin Plan. Prepared for the Northwest Power and Conservation Council. Prepared by the Yakama Nation, Klickitat County, and Washington Department of Fish and Wildlife.

Yakama Nation 2012. Draft Klickitat River Anadromous Fisheries Master Plan, Yakima/Klickitat Fisheries Program. Toppenish,WA.

Appendix 1- Take Tables

Estimated listed salmonid take levels by hatchery activity.

Steelhead

<i>ESU/Population</i>	Middle Columbia River Steelhead
<i>Activity</i>	Klickitat Hatchery Spring Chinook Program
<i>Location of hatchery activity</i>	Klickitat R. Hatchery
<i>Dates of activity</i>	Year-round
<i>Hatchery Program Operator</i>	YN & WDFW under co-managed YKFP

Type of Take	Annual Take of Listed Fish by life Stage (number of fish)			
	Egg/Fry	Juvenile/Smolt	Adult	Carcass
Observe or harass (a)		50-150	100	
Collect for transport (b)				
Capture, handle, and release (c)		500-2000 wild; 2000-3000 hatchery*	650-950 wild; 800-1000 hatchery**	
Capture, handle, tag/mark/tissue sample, and release (d)		3000-10,000 wild***		
Removal (e.g., broodstock) (e)				
Intentional lethal take (f)				
Unintentional lethal take (g)		190 wild; 150 hatchery	25 wild; 30 hatchery	
Other take (indirect, unintentional) (h)				

* Smolt trapping operations for monitoring purposes

** Although steelhead have not been taken during past hatchery practices, it is anticipated that adult steelhead will be collected and handled at the new collection facilities at Lyle Falls. Mortality occurs only on rare occasions during these operations.

***Stream fish (juvenile steelhead and resident trout) sampling operations

- a. Contact with listed fish through stream surveys, carcass and mark recovery projects, or migrational delay at weirs.
- b. Take associated with weir or trapping operations where listed fish are captured and transported for release.
- c. Take associated with weir or trapping operations where listed fish are captured, handled, and released upstream or downstream.
- d. Take occurring due to tagging and/or bio-sampling of fish collected through trapping operations prior to upstream or downstream release, or through carcass recovery programs.
- e. Listed fish removed from the wild and collected for use as broodstock.
- f. Intentional mortality of listed fish, usually as a result of spawning as broodstock.

- g. Unintentional mortality of listed fish, including loss of fish during transport or holding prior to spawning or prior to release into the wild, or, for integrated programs, mortalities during incubation and rearing.
- h. Other takes not identified above as a category

Take Table 2. Estimated listed salmonid take levels by hatchery activity.

Bull Trout

<i>ESU/Population</i>	Columbia River Basin DPS Bull Trout
<i>Activity</i>	Klickitat Hatchery Spring Chinook Program
<i>Location of hatchery activity</i>	Klickitat R. Hatchery
<i>Dates of activity</i>	Year-round
<i>Hatchery Program Operator</i>	YN & WDFW under co-managed YKFP

Type of Take	Annual Take of Listed Fish by life Stage (number of fish)			
	Egg/Fry	Juvenile/Smolt	Adult	Carcass
Observe or harass (a)		65*	15*	
Collect for transport (b)				
Capture, handle, and release (c)		5**	5**	
Capture, handle, tag/mark/tissue sample, and release (d)				
Removal (e.g., broodstock (e)				
Intentional lethal take (f)				
Unintentional lethal take (g)				
Other take (indirect, unintentional) (h)				

* Spawner surveys or habitat surveys

** Juvenile and adult trapping operations

- a. Contact with listed fish through stream surveys, carcass and mark recovery projects, or migrational delay at weirs.
- b. Take associated with weir or trapping operations where listed fish are captured and transported for release.
- c. Take associated with weir or trapping operations where listed fish are captured, handled, and released upstream or downstream.
- d. Take occurring due to tagging and/or bio-sampling of fish collected through trapping operations prior to upstream or downstream release, or through carcass recovery programs.
- e. Listed fish removed from the wild and collected for use as broodstock.
- f. Intentional mortality of listed fish, usually as a result of spawning as broodstock.
- g. Unintentional mortality of listed fish, including loss of fish during transport or holding prior to spawning or prior to release into the wild, or, for integrated programs, mortalities during incubation and rearing.
- h. Other takes not identified above as a category

YAKIMA/KLICKITAT FISHERIES PROJECT - KLICKITAT MONITORING AND EVALUATION PROJECT DESCRIPTION

Action	Overview of Action/Purpose	Description	Location	Anticipated Take (per year)				
				General ESA Approach/Notes	Bull Trout adults	Bull Trout juveniles	Steelhead adults	Steelhead juveniles
Spawning ground surveys	Document the abundance and temporal and spatial distribution of spawners and redds in the Klickitat basin for spring Chinook, fall Chinook, coho, and steelhead.	Conduct spawner surveys via wading and/or rafting within the known geographic range in the Klickitat subbasin. Count individual redds and record location using handheld GPS units. Record counts of live fish and carcasses. Collect biological information from carcasses (length, sex, scale sample, and tag/mark data). Examine carcasses for sex determination, egg/milt retention (percent spawned), and presence of decimal coded wire tags (CWT) tags or external experimental marks. Make attempts to cover the entire known spawning range of each species. Survey each stream reach multiple times (preferably at least 3 survey passes) during the spawning periods.	Throughout Klickitat subbasin (Klickitat River and anadromous-accessible tributaries); approximately 150 river miles of survey reaches	“Take” for this action may include scaring/stressing fish, temporary displacement of fish, and temporary interruption of spawning. Survey timing is as follows: spring Chinook - mid August through early October; fall Chinook - late October through early to mid December; coho - mid October through mid February; steelhead - late January through early June. No mortalities anticipated.	< 5	< 5	< 100	< 50
Adult salmonid monitoring at Lyle Falls Fishway	Collect data on adult salmonids in the Klickitat River to determine fish use, run timing, and estimate abundance.	Operate adult trap in the Lyle Falls fishway. Trap will be operated as flows and debris levels allow. Trap will be checked every 24 hours. Biological data will be collected including fish length, tag inspection, scale sample, and DNA samples. Marks (opercle punches and floy tags) will be administered to assist in subsequent resight/recapture and development of population estimates. Fish are crowded, then lifted and transported (via water-to-water transfer) to handling tanks for biological sampling. Fish are handled and sampled in small tanks using soft mesh knotless netting. Depending on funding, fish may also be PIT-tagged and/or radio-tagged to assist in determination of migration/holding patterns, spawning areas, passage issues, and fishway use. Appropriate anesthesia (generally electronarcosis) will be used; no chemical anesthetic is used due to harvest availability of fish after leaving trap.	RM 2.3 on the Klickitat River; T03N, R12E, Sec. 25 NWSW	“Take” for this action includes fish trapping and handling, collection of biological data and marking/tagging. Mortality only occurs on very rare occasions. Trap is operated year round as flows and debris/bedload levels allow.	< 5	0	600-800 Klickitat wild; 50 Snake R. wild; 800-1000 Klickitat hatchery; 50 Snake R. hatchery; < 3% mortality (Radio tagging: up to 100 wild, 100 hatchery of this total)	0
Adult salmonid monitoring at Castile Falls fishway	Collect data on adult salmonids in the Klickitat River to determine fish use, run timing, and estimate abundance	Operate enumeration facility in the Castile Falls fishway. Facility will be operated as flows and debris levels allow. Facility will be checked on a weekly or biweekly basis. Video and PIT tag detection equipment installed in fishway will be primary means of monitoring fish. Occasional trapping and handling of fish will occur for	RM 64.6 on the Klickitat River; T09N R13E, Sec. 18 SWSW	“Take” for this action includes occasional fish trapping and handling, collection of biological data and marking/tagging. Mortality will likely occur only on very rare occasions. Facility will be operated year round as flows and	0	0	50-100 Klickitat wild; < 2% mortality	0


		biological sampling purposes (fish length, tag inspection, scale sample, and DNA samples). When operated as a trap, facility will be checked every 24 hours. Handled fish will be placed in water-filled blackout tubes (PVC tubes with cutout sections) or soft mesh knotless netting; appropriate anesthesia (e.g., electronarcosis or MS-222) will be used.		debris/bedload levels allow.				
Juvenile outmigration monitoring	Continuous monitoring of juvenile outmigration in the upper and lower Klickitat River utilizing rotary screw traps. Information to provide an index of number of smolts, parr, and fry leaving the Klickitat system.	Operate floating rotary screw traps to monitor juvenile (smolt, parr, and fry) outmigration in the upper and lower Klickitat River. Traps will be fished year round (as flows, debris levels, and hatchery releases allow) at the: 1. Lyle Falls trap, 2. Klickitat Hatchery trap, and 3. seasonally (between May and November) at the Castile Falls trap. Calibration studies (mark-recapture trials) will be conducted to estimate trap efficiency and assist in development of smolt production estimates. Environmental and trap data will be recorded along with bio-data on 10 to 30 of each salmonid species represented. Fish will be anesthetized and sampled for length, weight, scales, and DNA. Additional tags or marks may also be administered (fin clips for mark-recapture efficiency testing and PIT tags). The excess and non-salmonid fish will be tallied by species. Depending on funding, 1 or 2 additional smolt traps or instream PIT tag antennas may temporarily be deployed in selected key tributaries (e.g. White Creek).	Lyle Falls (RM 2.3 on the Klickitat River; T03N,R12E, Sec. 25 NWSW) Klickitat Hatchery (RM 42 on the Klickitat River; T06N R13E, Sec. 4 SWNE) Castile Falls (RM 64 on the Klickitat River; T09N,R13E, Sec.19 NENE)	“Take” for this action includes fish trapping and handling, collection of biological data and marking/tagging. Mortality only occurs on rare occasions. Lyle Falls and Klickitat Hatchery traps are operated year round (as flows, debris levels, and hatchery releases allow); Castile Falls trap is operated seasonally (generally May to November)	< 5	< 5	< 5 (kelts)	500-2000 wild; 2000-3000 hatchery; < 5% mortality (PIT tagging: up to 1500 wild of this total; PIT tagging may also include an additional 10,000 hatchery smolts tagged at Skamania Hatchery)
Juvenile and resident salmonid population surveys	Determine the spatial distribution, abundance, survival, and migration patterns of salmonids throughout the basin to provide baseline information and evaluate hatchery and habitat actions.	Electrofishing surveys will be conducted in selected key tributary and mainstream reaches. Standard depletion or mark-recapture estimates to determine abundance will be utilized. Snorkel surveys will also be used in selected reaches. Population surveys may be completed in selected reaches pre and post-habitat improvement actions. The number of sites sampled will be determined by time allotted to other field season activities (e.g., habitat surveys). Fish captured will be anesthetized and sampled for length, weight, scales, and DNA. Depending on funding, additional tags or marks may also be administered (e.g., fin clips and PIT tags for survival estimation and juvenile outmigration monitoring).	Throughout Klickitat subbasin; includes White Cr. watershed, other specific locations to be determined	“Take” for this action includes fish capture (via electrofishing), scaring/stressing fish or temporary displacement of fish (during snorkel surveys), handling, tagging, and collection of biological data. Mortality only occurs on very rare occasions. Sampling would generally occur in the summer and early fall.	< 5	< 5	< 5	3000-10000 wild; <3% mortality (PIT tagging: up to 5000 wild of this total)
Scale analysis	Scales are taken at traps and from carcasses encountered on spawner surveys as part of a	Fish scales are taken at screw traps, at the Lyle and Castile adult traps, and from carcasses encountered on spawning surveys. The majority of the scale reading is done by YKFP M&E staff; some scales may be read by WDFW staff.	Same location as screw traps, adult traps, and spawner surveys (above)	No effect – action is scale reading and analysis;	No effect – action is scale reading and analysis	No effect – action is scale reading and analysis	No effect – action is scale reading and analysis	No effect – action is scale reading and analysis

	continuous and ongoing sampling routine to determine age and stock composition of juvenile and adult salmonid stocks in the Klickitat basin.							
Sediment monitoring	Monitor stream sediment loads associated with natural and anthropogenic factors (e.g., logging, agriculture, and road building) which can increase sediment loads in streams utilized by all salmonids in the Klickitat basin.	Gravel samples will be collected and analyzed using Washington State DNR Timber, Fish and Wildlife (TFW) monitoring methodology. McNeil gravel core samples will be collected at 10-12 sites and will be sieved to estimate percentage composition of various substrate particle sizes.	Klickitat River between RM 16 and 88; Diamond Fork Cr. between RM 0 and 12; White Cr. RM 9; Tepee Cr. RM 5	“Take” for this action may include scaring/stressing fish, temporary displacement of fish, and minor localized turbidity increases during gravel sampling. Sampling would occur in the fall (October – November). No anticipated mortalities.	0	0	<5	<10
Water quality monitoring	Continue ongoing water quality monitoring at established and selected new sites.	Record water quality measurements on selected tributaries and within selected habitat survey reaches on a seasonal and as-possible basis. Portable field meters will be used to measure and record the following parameters: temperature, dissolved oxygen, pH, conductivity, and turbidity. Data will be recorded at 36-38 locations, approximately 5-8 times per year at each location. Temperature is also continuously monitored via data loggers placed in streams at these locations.	Approximately 38 locations throughout Klickitat subbasin (Klickitat R. and tributaries)	No effect is anticipated.				
Habitat Surveys	Complete habitat surveys at selected sites and reaches. Quantitative habitat data will provide the foundation for decision-making relative to habitat restoration, as well as refining related attributes of the EDT model. Survey data will also assist in effectiveness monitoring of	Collect baseline data on existing habitat conditions throughout the basin. The habitat inventories will be conducted using YKFP Rapid Habitat Assessment protocols and the Washington State DNR Timber, Fish and Wildlife (TFW) monitoring methodology (modules: Stream Segment Identification, Reference Point Survey, Habitat Unit Survey, and Large Woody Debris Survey). Sites may include previously surveyed sites and/or new sites.	Throughout Klickitat subbasin; specific locations to be determined	“Take” for this action may include scaring/stressing fish or temporary displacement of fish during habitat survey. Surveys generally occur in late spring and summer. No anticipated mortalities.	0	< 50	0	< 100

	habitat restoration projects and in other land management planning.							
Genetic data collection, analysis, and synthesis	Provide information on subpopulation structure, geographic variation, and production in order to minimize any effects from hatchery actions.	Genetic samples will be collected from adult salmonids at the Lyle and Castile adult traps, and from juveniles at rotary screw traps. Additional samples may also be collected via stream electrofishing. Samples will be sent to Columbia River Intertribal Fish commission (CRITFC) genetics lab or other genetics labs for analysis. YKFP biologists, in collaboration with CRITFC geneticists, will compile existing data and analyze genetics information. May also include research on phenotypic expression of biologically functional genes (e.g. thermal tolerance and smoltification).	Same location as screw traps, adult traps, and spawner surveys (above) Other locations to be determined throughout Klickitat subbasin.	Take for this action includes fish capture and handling (with non-lethal fin clip, fin punch, or opercle punch sampling) during stream electrofishing and adult trap and screw trap operation. May include some lethal sampling (up to 100 juvenile steelhead per year) for research on phenotypic expression of biologically functional genes; otherwise no mortalities expected.	0	0	0	500-1000 (most of this total included as part of other sampling described above)

Appendix 2 - Steelhead and Bull Trout Distribution

Steelhead Distribution



Bull Trout Distribution

